

ANTIMICROBIAL CONSTITUENTS FROM *PLUCHEA WALLICHIANA* DC.

Mehdi Hassan Kazmi, Sadia Khan and Saira Hameed

Department of Applied Chemistry and Chemical Technology, University of Karachi, Karachi- 75270, Pakistan

ABSTRACT

Eight compounds viz., β -amyrin (**1**), β -amyrin acetate (**2**), (+)-syringaresinol (**3**), *ent*-pimara-8(14), 15-dien-19-oic acid (**4**), pluviatilol (**5**), β -sitosterol (**6**), β -sitosterol 3-*O*- β -D-glucopyranoside (**7**), and apigenin 7-*O*- β -D-glucoside (**8**), were isolated for the first time from *Pluchea wallichiana* DC. and their structures were elucidated by extensive spectroscopic studies. All the isolated compounds were tested for their antimicrobial activity.

Keywords: *Pluchea wallichiana*, antimicrobial activity, phytochemistry

INTRODUCTION

The genus *Pluchea* belongs to the Family Compositae which is widely distributed in Pakistan, India, Afghanistan and Iran (Mhaskar *et al.*, 2002; Kupchan *et al.*, 1970). The family Compositae contains Sesquiterpene, Chlorogenic acid, lactones, orange yellow pigment, sitosterol glycoside, which are known to possess cytotoxic, antitumor and mutagenic properties (Therdore, 1908; Perry and Metzgar, 1980; Jayawera, 1980). Several species have shown antifungal, antibacterial and anti-inflammatory activities against number of microorganisms. The plants are traditionally used in the treatment of peptic, ulcer burns, abdominal pain and bacterial diseases. (Jafri, 1966; Ahmed and Fiza, 1988). The literature survey reveals that there are no phytochemical work has so far been carried out on *Pluchea wallichiana* DC. This led us to carry out phytochemical investigations on this plant.

MATERIAL AND METHODS

Plant Material:

The whole plant of *Pluchea wallichiana* DC. was collected by botanical authority from Karachi region in the month of July to August and identified by Dr. Surraya Khatoon, Plant Taxonomist, Department of Botany University of Karachi, Pakistan, where a voucher specimen (1037/KUH) has been deposited.

Isolation:

The shade dried whole plants of *P. wallichiana* (20Kg) was extracted with MeOH (3x50L) at room temperature. The combined methanolic extract was evaporated under reduced pressure to obtain a thick gummy mass (700gm). It was suspended in water and successively extracted with *n*-hexane, ethylacetate and *n*-butanol. The chloroform soluble fraction was subjected to column chromatography over silica gel eluting with *n*-hexane-chloroform, chloroform and chloroform-methanol in increasing order of polarity to obtain six major fractions C_A-C_F. The fraction C_A obtained from *n*-hexane-chloroform (6.0: 3.0) gave two spots on TLC which on preparative TLC in the same solvent system gave β -amyrine (**1**) (30 mg) and β -amyrine acetate (**2**) (20 mg) respectively. The fraction C_B obtained from the *n*-hexane-chloroform (5.0 : 5.0) was rechromatographed over silica gel with the same solvent and final purification with PTLC using *n*-hexane-chloroform (4.5 : 5.5) as solvent system provided (+)-syringaresinol (**3**) (20 mg). The fraction C_C obtained from *n*-hexane-chloroform (4.0: 6.0) were combined and rechromatographed over silica gel eluting with same solvent system. The final purification was achieved through PTLC using *n*-hexane-chloroform (3.0 : 7.0) as eluent to afford *ent*-pimara-8-(14),-15-dien-19-oic acid (**4**) (20 mg). The fraction C_D obtained from *n*-hexane-chloroform (2.0: 8.0) gave two spots on TLC which on PTLC with the same solvent system gave pluviatilol (**5**) (25 mg) and β -sitosterol (**6**) (30 mg). The fraction C_E obtained from chloroform-methanol (9.8: 0.2) gave one major spot on TLC which on PTLC in the solvent system chloroform-methanol (9.5:0.5) gave β -sitosterol 3-*O*- β -D-glucopyranoside (**7**) (25 mg). The fraction C_F obtained from chloroform-methanol (9.5 : 0.5) were combined and rechromatographed over silica gel eluting with the solvent system (9.4 : 0.6) and final purification from PTLC using chloroform-methanol (9.2 : 0.8) as developing solvent yielded apigenin 7-*O*- β -D-glucopyranoside (**8**) (25 mg). All the compounds were identified through comparison of physical and spectral data with those reported in literature.

β -Amyrin (1)

Crystallization from ethanol (30 mg), m.P 197-198°C; $[\alpha]_D^{25}$: +100° ($c = 0.21$, CHCl₃). IR (KBr) ν_{max} cm⁻¹: 3510 (hydroxyl), 3055, 1635 and 820. ¹H-NMR (CDCl₃, 500 MHz): δ 5.11 (1H, m, H-12), 3.20 (1H, dd, $J = 10.0, 4.5$ Hz, H-3), 1.02, 1.01, 1.08, 0.96, 0.93, 0.88, 0.85 and 0.80 (3H, each s, Me). HREIMS m/z : 426.3825 (calcd. for C₃₀H₅₀O, 426.3861). The physical and spectral data showed complete agreement with those reported in the literature (Budzikiewicz *et al.* 1963 ; Shamma *et al.*, 1962).

 β -Amyrin acetate (2)

Crystallization from methanol (20 mg), m.P 244-245°C; $[\alpha]_D^{25}$: + 81.4° ($c = 0.20$, CHCl₃). IR (KBr) ν_{max} cm⁻¹: 3055, 1710, 1660, 1460, 1382, 1180 and 810 cm⁻¹. ¹H-NMR (CDCl₃, 500 MHz): δ 5.10 (1H, m, H-12), 4.05 (1H, dd, $J = 10.0, 4.5$ Hz, H-3), 2.13 (3H, s, OAc), 1.00, 1.07, 0.98, 0.96 (3H, each s, Me), 0.90 (6H, s, H-29 and H-30), 0.86 and 0.81 (3H, each s, Me). HREIMS m/z : 468.3931 (calcd. for C₃₂H₅₂O₂, 468.3916). The physical and spectral data corresponded to the reported values (Chow and Quon, 1970).

(+)-Syringaresinol (3)

Crystallization from methanol (20 mg), m.P 179-180°C; $[\alpha]_D^{20}$: + 6.9 (C 0.02, CHCl₃). IR (KBr) ν_{max} cm⁻¹ 3428, 1614, 1520. ¹H-NMR (CDCl₃, 500 MHz): δ 3.11 (1H, m, H-1), 4.73 (1H, d, $J = 4.5$ Hz, H-2), 3.89 (1H, m, H-4), 4.30 (1H, dd, $J = 7.0, 9.0$ Hz, H-4), 3.12 (1H, m, H-5), 4.73 (1H, d, $J = 4.5$ Hz, H-6), 3.89 (1H, m, H-8), 4.30 (1H, dd, $J = 7.0, 9.0$ Hz, H-8), 6.58~6.60 (4H, H-2',6',2'',6''), 3.90, (3H, s, OCH₃) HREIMS m/z 418. 4269 (calcd, for C₂₂H₂₆O₈, 418.1627). The physical and spectral data were in agreement with those reported in literature (Deyama *et al.*, 1987; Macrae and Towers, 1985; Tanaka *et al.*, 1989).

ent-Pimara-8(14), 15-dien-19-oic acid (4)









Crystallization from methanol (30 mg), m.P 165-166°C; IR ν_{max} (KBr) cm⁻¹, 3400, 1690, 1460 ¹H-NMR (CDCl₃, 300 MHz): δ 0.66 (3H, s, H-20), 1.01 (3H, s, H-17), 1.23 (3H, s, H-18), 4.91 (1H, dd, $J = 2.1, 11.7$ Hz, H-16a), 4.96 (1H, dd, $J = 1.8, 5.1$ Hz, H-16b), 5.16 (1H, s, H-14), 5.71 (1H, dd, $J = 10.5, 17.1$ Hz, H-15). HREIMS m/z 302.4391 (calcd, for C₂₀H₃₂O₂, 302.4421). The physical and spectral data showed complete agreement with those reported in the literature (Sy and Brown, 1998).

Pluviatilol (5)

Crystallized with benzene and petroleum ether (25 mg), m.P 161-163°C; $[\alpha]_D^{20}$ + 79.8 (C 0.16 MeOH). ¹H-NMR (CDCl₃, 500 MHz): δ 2.91 (1H, m, H-1), 4.42 (1H, d, $J = 7.5$ Hz, H-2), 3.32 (1H, m, H-4), 3.85 (1H, m, H-4), 3.32 (1H, m, H-5), 4.86 (1H, d, $J = 6.0$ Hz, H-6), 3.85 (1H, dd, $J = 6.5, 9.5$ Hz, H-8), 4.13 (1H, dd, $J = 1.0, 9.5$ Hz, H - 8), 6.81~6.89 (6H, H-2',5',6',2'',5'',6''), 5.97, (2H, s, OCH₂O), 3.91 (3H, s, OCH₃). HREIMS m/z : 356.3740 (calcd. for C₂₀H₂₀O₆, 356.1259). The physical and spectral data corresponded to the reported values (Corrie *et al.*, 1970; Banerji and Pal, 1982).

 β -Sitosterol (6)

Crystallization from acetone (30 mg), m.P 135°C; $[\alpha]_D^{25}$: + 35.5 ($c = 0.22$, CHCl₃). IR (KBr) ν_{max} : 3446, 3050, 1650 and 815 cm⁻¹. ¹H-NMR (CDCl₃, 400 MHz) δ : 5.11 (1H, m, H-6), 3.36 (1H, m, H-3), 1.01 (3H, s, Me-19), 0.92 (3H, d, $J = 6.2$ Hz, Me-21), 0.84 (3H, t, $J = 7.0$ Hz, Me-29), 0.83 (3H, d, $J = 6.5$ Hz, Me-26), 0.81 (3H, d, $J = 6.5$ Hz, Me-27), 0.68 (3H, s, Me-18). HREIMS: m/z 414.3845 (calcd. for C₂₉H₅₀O, 414.3861). The physical and spectral data showed complete resemblance with those reported in the literature (Holland *et al.*, 1978; Rubintein *et al.*, 1976).

***β*-Amyrin (1)****1****(+)-Syringaresinol (3)****3*****β*-Amyrin acetate (2)****2****ent-Pimara-8(14), 15-dien-19-oic acid (4)****4*****β*-Sitosterol (6)****6****Pluviatilol (5)****5*****β*-Sitosterol 3-*O*-*β*-D-glucopyranoside (7)****7****Apigenine 7-*O*-*β*-D-glucoside (8)****8**

β -Sitosterol 3-O- β -D-glucopyranoside (7)

Crystallization from methanol (25 mg), m.P 279-280 °C; $[\alpha]_D^{25}$ -14.5 ($c = 0.3$, MeOH) IR (KBr) ν_{max} : 3452, 3044, 1646, 1618, 1595-1550 cm^{-1} . 1H -NMR ($CDCl_3$, 300 MHz) δ : 5.33 (1H, d, $J = 7.2$ Hz, H-1'), 5.12 (1H, br d, $J = 5.4$ Hz, H-6), 3.85 (1H, m, H-3), 3.82-4.42 (m, Glc-H), 1.01 (3H, s, Me-19), 0.92 (3H, d, $J = 6.2$ Hz, Me-21), 0.84 (3H, t, $J = 7.0$ Hz, Me-29), 0.83 (3H, d, $J = 6.5$ Hz, Me-26), 0.81 (3H, d, $J = 6.5$ Hz, Me-27), 0.68 (3H, s, Me-18). HRFABMS: m/z 576.4386 (calcd for $C_{35}H_{61}O_6$ 576.43897). The physical and spectral data corresponded to the reported values. (Iribarren *et al.*, 1983; Wyllie *et al.*, 1976).

Apigenine 7-O- β -D-glucoside (8)

Amorphous powder (25 mg), $[\alpha]_D^{25}$: -64.0 ($c = 0.14$, MeOH). UV λ_{max} (log ϵ) (MeOH) nm: 6.9 (4.42) and 269 (3.03). IR (KBr) ν_{max} cm^{-1} : 3370 and 1660. 1H -NMR (CD_3OD , 400 MHz) δ : 7.86 (2H, d, $J = 9.0$ Hz, H-2' and H-6'), 6.92 (2H, d, $J = 9.0$ Hz, H-3' and H-5'), 6.75 (1H, d, $J = 3.0$ Hz, H-8), 6.68 (1H, s, H-3), 6.43 (1H, d, $J = 3.0$ Hz, H-6), 5.04 (1H, d, $J = 7.5$ Hz, H-1'), 4.52 (1H, m, H-4'), 3.84 (1H, m, H-2'), 3.78 (2H, m, H-6''), 3.59 (1H, m, H-3'') and 3.38 (1H, m, H-5').

HRFABMS (+ve) $[M+H]^+$ m/z : 433.1135 (calcd. for $C_{21}H_{20}O_{10}$, 433.1125). The physical and spectral data showed complete resemblance with those reported in literature (Sadikum *et al.*, 1980; Redacchi *et al.*, 1980).

Antimicrobial Assay:

The antibacterial and antifungal were determined by agar diffusion method proposed by (Jaffer *et al.*, 1988), one loop full of 24hr- old culture of selected bacteria Spread on the surface of Mueller Hinton Agar Plates wells were drug in the medium with the help of sterile borer. A stock solution of extract 2mg/ml was prepared in DMSO and dilution of the stock solution containing hundred microlitre of each dilution was added to their respective wells and after 24hr-growth inhibition of bacteria was observed.

Table 1. Antibacterial and antifungal activity of compounds 1-8 from *Pluchea wallichiana*.

Microorganism	Zone of inhibition (mm)								
	Compounds (1-8)								
Bacteria	1	2	3	4	5	6	7	8	Control
Gram positive									
<i>Bacillus subtilis</i>	10	15	28	20	5	0	20	0	+
<i>Shigella flenneri</i>	15	20	22	15	15	15	25	25	+
<i>Staphylococcus aureus</i>	5	20	16	12	15	12	18	20	+
<i>Streptococcus pyogens</i>	12	5	12	10	10	20	20	15	+
<i>Streptococcus agalactae</i>	12	0	20	13	10	18	20	20	+
Gram negative									
<i>Escherichia coli</i>	15	18	20	18	7	5	22	15	+
<i>Pseudomonas aeruginosa</i>	12	0	20	17	12	3	18	0	+
<i>Salmonella paratyphi</i>	10	15	25	15	12	4	16	16	+
<i>Shigella bydi</i>	18	16	15	10	10	6	15	14	+
<i>Salmonella typhi</i>	20	12	12	15	8	2	15	12	+
Fungi									
<i>Aspergillus nigar</i>	2	0	58	25	7	12	15	16	+
<i>A. fumigatus</i>	0	10	30	12	5	10	20	20	+
<i>A. flavus</i>	5	12	20	17	0	10	17	25	+
<i>Alternaria solani</i>	7	14	25	15	5	5	12	17	+

RESULTS AND DISCUSSION

The methanolic extract was fractionated into n-hexane, chloroform, ethyl acetate, n-butanol and water soluble fractions. Of these chloroform fraction has been studied in the present investigations. A series of column chromatographic techniques resulted in isolation of eight known compounds reported for the first time from this species. These could be identified as β -amyirin (**1**), β -amyirin acetate (**2**), (+) – syringaresinol (**3**), *ent*-pimara- 8 (14), 15-dien-19-oic acid (**4**), pluviatilol (**5**), β -sitosterol (**6**), β -sitosterol 3-*O*- β -D – glucopyranoside (**7**), and apigenin 7-*O*- β -D-glucoside (**8**). The isolated compounds were tested for their antimicrobial activity (Table 1). The compounds 3, 4, 7 and 8 showed significant antimicrobial activity against *Bacillus subtilis*, *Shigella flenneri*, *Staphylococcus aureus*, *Streptococcus pyogens*, *Streptococcus agalactae*. Other compounds showed weak to moderate activity.

REFERENCES

- Ahmed, V. U. and K. Fiza (1988). Structure of a new Eudesmane sesquiterpene Argutinin from *Pluchea arguta* Boiss. *Naturforsch B Chem Sci.*, 43 (7): 911.
- Ali, S. I. and M. Qaiser (2003). *Flora of Pakistan*. Missouri Botanical Press, Missouri Botanical Garden st. Louis, Missouri, U. S. A., p103, 105.
- Banerji, A. and S. Pal (1982). Constituents of *Piper sylvaticum*: Structure of Sylvatesmin. *J. Nat-Prod.*, 45: 672.
- Budzikiewicz, H., J.M. Wilson and C. Djerassi (1963). Mass spectrometry in structural and stereochemical problems, Pentacyclic Triterpenes. *J. Am.Chem. Soc.*, 85: 3688.
- Chow, Y. L. and H.H. Quon (1970). Biogenetically related triterpene from *Elateriospermum Tapos* Bark. *Phytochemistry*, 9: 1151.
- Compodium (2002). Medicinal plants used in Malaysia II, Herbal Medicinal Research Centre Institute for Medical Research Kuala Lumpur Malaysia, p 239.
- Corrie, J. E. T., G. H. Green and W. C. Taylor (1970). The chemical constituents of Australian *Zanthoxylum* species. V. The constituents of *Z. Pluviatile* Hartley, the structures of two new lignans. *Aust. J. Chem.*, 23: 133 .
- Deyama T., T. Ikawa, S. Kitagawa and S. Nishibe (1987). The constituents of *Eucommia OLIV.V*. Isolation of Dihydroxydehydrodiconiferyl alcohol isomers and phenolic compounds. *Chem. Pharm. Bull.*, 35: 1785.
- Holland H. L., P. R. P. Diakow and G. J. Taylor (1978). ¹³C nuclear magnetic resonance spectra of some C-19-hydroxy, C-5,6 epoxy, C-24 ethyl, and C- 19- norsteroids. *Can. J. Chem*, 56: 3121..
- Iribarren, A. M. and A.B. Pomilio (1983). Sitosterol 3- 0- α -D- Xyluronofuranoside from *Bauhinia candicans*. *J. Nat. Prod.*, 46: 725.
- Jaffer, H.J., A.L.M. Jawad, H.S. Saber and A. Al-Naib (1988). Evaluation of antimicrobial activity of *Withania somnifera* extracts. *Fitoterapia*, 59: 497.
- Jafri S. M. H. (1966). *The flora of Karachi*. Book Corporation, Karachi. p 329.
- Jayawera, D. M. A. (1980). *Medicinal plants indigenous and Exotic* Vol II. National Science Council of Sri Lanka Colombo, p 45.
- Krishna, A. (2003). *The wealth of India*. Council of Scientific and Industrial Research New Delhi VIII,p 161,
- Kupchan S.M., D. C. Fessler, M .A. Eakin and T..J. Giacobba (1970). Reactions of alpha methylene lactone tumour inhibitors with model biological nucleophiles. *Science*, 168: 376.
- Macrae W. D., and G. H. N. Towers (1985). Non alkaloidal constituents of *Virola elongata* bark. *Phytochemistry*, 24: 561.
- Mhaskar K. S., E. Blatter and J.F. Caius (2002). *Indian Medicinal plants*. Sri Satguru Publications Delhi, 6, p. 1817 .
- Perry L. M. and J. Metzger (1980). *Medicinal Plants of East and South Asia*. The MIT Press Cambridge England, 96, p97.
- Redacelli, C., L. Formentini and E. Santaniello (1980). Apigenin 7- Glucoside and its 2"- and 6" – acetates from Ligulate flowers of *Mmatricaria Chamomilla*. *Phytochemistry*, 19: 985 .
- Rubinstein, I., L. J. Goad, A. D. H. Clague and J. L. Mulheirn (1976). The 220 MHz NMR spectra of phytosterols. *Phytochemistry*, 15: 195.
- Sadikum, A., I. Aninah, N. Ismail and P. Ibrahim (1980). Sterols and sterol glycosides from the leaves of *Gynura procumbens*. *Nat. Prod. Sci.*, 2: 19 .
- Shamma M., R. E. Glick and R.O. Mumma (1962). The nuclear magneic resonance spectra of pentacyclic triterpenes, *J. Org. Chem.*, 27: 4512.
- Sy, L. K. and G. D. Brown (1998). Abietane Diterpenes from *Illicium Angustisepalum*. *J. Nat. Prod.*, 61: 907.

- Tanaka, H., T. Nakamura, K. Ichino and K. A. Ito (1989). Lignan from *Actinodaphne Longifolia*. *Phytochemistry*, 28: 952.
- Therdore C. I. E. C. (1908). *Flora of the presidency of Bombay II*. Under the Authority of the secretary of Estate for Indica in Council, p.1.
- Wyllie, S. G., B. A. Amos and L. Tokes (1976). Electron impact induced fragmentation of cholesterol and related C-5 unsaturated steroids. *J. Org. Chem.*, 42: p725.

(Accepted for publication July 2012)