

## CHARACTERIZATION OF INVERTASE AND ALPHA AMYLASE FROM TWO FUNGAL SPECIES, *PENICILLIUM LILACINUM* AND *ASPERGILLUS NIGER*

Kashif Ahmed\*, Shamsa Naz and Anjum Ayub

Department of Chemistry, NED University of Engineering and Technology, Karachi, Pakistan

\*Corresponding author's E-mail: [kashifahm@neduet.edu.pk](mailto:kashifahm@neduet.edu.pk), [kashif25473@yahoo.com](mailto:kashif25473@yahoo.com)

---

### ABSTRACT

Amylases are one of the key enzymes that are used in food and other biotechnological applications. These enzymes hydrolyze starch into polymers composed of glucose units and have prospective application in food, fermentation and pharmaceutical industries. Although Amylases can be obtained from sources such as plants, animals and microorganisms, the enzymes from fungal and bacterial species show profound applications in industries. We have identified two fungal species, *Penicillium lilacinum* and *Aspergillus niger* that efficiently produce invertase and  $\alpha$ -amylase. We have also identified optimal conditions for their production. The highest quantity of invertase (13.05 U/mL) was obtained from *Penicillium lilacinum* under these conditions: CM1, culture medium; yeast extract, nitrogen source; date syrup, carbon source; 96 h, incubation time period; culture medium pH 8.0; 40°C, incubation temperature 40°C; inoculum size conidia  $6 \times 10^6$ , agitation rate 200 rev/min. The highest amount of  $\alpha$ -amylase (8.14 U/mL) was obtained by *Aspergillus niger* under conditions: M1, culture medium; yeast extract as, nitrogen source; molasses as carbon source; incubation time period 72 h; initial pH of culture medium 6.5; incubation temperature 40°C; inoculum size  $5 \times 10^6$  conidia; agitation rate 150 rev/min. These strains are potential candidates for industrial use because these enzymes maintain their activities even at harsh pH and temperature conditions.

**Keywords:**  $\alpha$ -amylase, Commercial enzyme, Industrial enzyme, Invertase, Submerged fermentation.

---

### INTRODUCTION

Alpha-amylase or 1, 4-alpha-D-glucan glucanohydrolase, E. C. 3.2.1.1, is an extracellular enzyme, which splits  $\alpha$ -1, 4- glycosidic bonds of starch (a polysaccharide and composed of amylopectin and amylose) and produces alpha limit dextrin, maltose and glucose (Ahmed *et al.*, 2015b; 2015c). Invertase and amylase have been extensively used commercial enzymes in medicinal, clinical, food and analytical chemistry and also in the brewing, baking, paper, pharmaceutical, detergent and textile industries (Ahmed *et al.*, 2015b; 2015c; Sundarram and Murthy, 2014).

In many countries agricultural wastes are usually burnt in open air that can add to air pollution. By using agricultural (Cellulosic) wastes as energy source at least two targets can be achieved; on one hand air pollution can be contained and at the same time useful products can be produced. A body of literatures has reported nonconventional energy sources such as cassava starch, date syrup, starch, cotton stalk, rice husk, wheat straw, potato peel, sunflower waste, oilcakes, tapioca, fruit peel, corn and many others have been used in the fermentation process for enzymes production (Ahmed *et al.*, 2015a, 2015b).

At the moment the biggest commercial application of amylases are in food industries. The biochemical multiplicity of microorganisms makes them reasonable sources of a varied selection of enzymes for use in food and other biotechnological usage (Mamma *et al.*, 2008; Ahmed *et al.*, 2015b; 2015c; Kulshrestha, 2013; Ahmed *et al.*, 2015b; 2015c). We have investigated fungal  $\alpha$ -amylases from *Penicillium lilacinum* and *Aspergillus niger*, physical and chemical constraint, and the use of these enzymes in industrial and other applications. These fungal species were grown on various agricultural based cellulosic wastes.

### MATERIALS AND METHODS

#### Optimization of Enzyme Production Parameters:

All experiments were done in such a way that the parameter optimized in one experiment was fixed in the subsequent experiments for the maximum production of enzyme. Following were parameters:

**Culture media:** First of all the most suitable culture medium was determined. For optimization of  $\alpha$ -Amylase production following culture media were used having composition (g/L).

**M1:** Dextrose 10, Peptone 5, Epsom salt 5,  $\text{KH}_2\text{PO}_4$  5, Common salt 2.5, ferrous sulphate hepta hydrate .01,  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  0.002,  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$  0.001 and thiamine hydrochloride 0.001 (Burrel *et al.*, 1966).

**M2:** Soluble starch 20,  $\text{NH}_4\text{NO}_3$  10,  $\text{KH}_2\text{PO}_4$ , 14, KCl, 0.5, Epsom salt 0.1,  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.01 (Matthias, 2013).

**M3:** NaCl 0.8, KCl 0.8, CaCl<sub>2</sub> 0.1, Na<sub>2</sub>HPO<sub>4</sub> 2.0, MgSO<sub>4</sub> 0.2, FeSO<sub>4</sub> 0.1, 8.0 Glucose, NH<sub>4</sub>Cl 2.0 (Khan and Yadav, 2011).

**M4:** ZnSO<sub>4</sub>.7H<sub>2</sub>O 0.062, FeSO<sub>4</sub> 0.068, copper sulphate pent hydrate 0.0001 and wheat bran 100 (Hayashida and Teramoto, 1986).

For invertase following were composition of culture media

**CM1:** Dextrose 10, peptone 5, Epsom salt 5, KH<sub>2</sub> PO<sub>4</sub> 5, common salt 2.5, ferrous sulphate hepta hydrate 0.01, ZnSO<sub>4</sub>.7H<sub>2</sub>O 0.002, MnSO<sub>4</sub>.H<sub>2</sub>O 0.001 and thiamine hydrochloride 0.001 (Burrel *et al.*, 1966).

**CM2:** Yeast extract 10, peptone 20 and sucrose 20 (Dworschock and Wickerham, 1961).

**CM3:** Yeast extract 20, peptone 40, sucrose 20, KH<sub>2</sub>(PO<sub>4</sub>)<sub>2</sub> and Epsom salt 1 (Souza *et al.*, 2007).

**CM4:** NaNO<sub>3</sub> 3, KCl 0.5, Epsom salt 0.5, ferrous sulphate hepta hydrate 0.01, K<sub>2</sub>HPO<sub>4</sub> 1, Sucrose 30 (Almeida *et al.*, 2005).

**CM5:** Sucrose 40, corn steep liquor 30, NaNO<sub>3</sub> 3, KH<sub>2</sub>PO<sub>4</sub> 0.5, Epsom salt 0.05, CaCO<sub>3</sub> 2.5 (Poonawalla *et al.*, 1965).

After the selection of the most suitable culture medium carbon source, nitrogen source, initial pH of culture medium, incubation temperature, conidia count and agitation rate were checked for the maximum production of enzymes in a sequence (Ahmed *et al.*, 2015a, 2015b)

### Invertase activity

Invertase activity was determined as described by Ahmed *et al.* (2015a; 2016). In brief, 0.1 mL of enzyme sample was mixed with 2.5 mL acetate buffer (50 mM, pH 5.5) and 0.1 mL 300 mM of sucrose; then incubated for 5 minutes at 35° C, then added 1.0 mL of dinitro Salicylic acid. All contents then boiled for five minutes and noted Absorbance at 540 nm.

### $\alpha$ -Amylase activity

$\alpha$ - Amylase Activity was determined as described by Ahmed *et al.* (2015b; 2015c). In brief, 1.0 mL of enzyme sample is mixed with 1.0 mL of soluble starch (1 % w/v) in 50 mM sodium phosphate buffer at pH 7.0 and then incubated for 5 min at 50° C, then added 1.0 mL dinitro Salicylic acid and then boiled for five minutes and noted Absorbance at 540 nm.

Table 1. Optimal Conditions for the production of invertase by various fungi.

Fungi	Culture medium	Incubation time period	Carbon source	Nitrogen source	Incubation Temperature	Initial pH	Inoculum Size	Agitation Rate	Optimized invertase activity
		h			° C		Conidia/mL	Rev/min	U/mL
<i>A. niger</i>	CM1	72	Molasses	yeast extract	40	6.0	5x10 <sup>6</sup>	150	8.23
<i>A. fumigatus</i>	CM1	48	Sunflower waste	yeast extract	30	6.5	6x10 <sup>6</sup>	100	6.37
<i>P. notatum</i>	CM1	48	Molasses	yeast extract	40	6.5	6x10 <sup>6</sup>	200	6.41
<i>P. lilacinum</i>	CM1	96	date syrup	yeast extract	40	8.0	6x10 <sup>6</sup>	200	13.05
<i>M. geophillus</i>	CM1	48	molasses	yeast extract	35	6.5	5x10 <sup>6</sup>	150	5.25
<i>P. expansum</i>	CM1	48	date syrup	yeast extract	35	5.0	6x10 <sup>6</sup>	150	6.37

## RESULTS

### Determine optimal conditions for invertase production

We obtained the maximum quantity of invertase (13.05 U/mL) was obtained by *Penicillium lilacinum* (Table 1) under conditions: CM1, culture medium; yeast extract as nitrogen source; date syrup as, carbon source ; 96 h of

incubation time period; 8, initial pH of culture medium; 40° C, incubation temperature;  $6 \times 10^6$  conidia, inoculum size; 200 rev/min, agitation rate. Optimal conditions for the maximum production of invertase by other fungi were *Aspergillus fumigatus*, 8.23 U/mL; *Aspergillus niger*, 6.24 U/mL; *Penicillium notatum*, 6.41 U/mL; *Penicillium lilacinum*, 13.05 U/mL; *Mucor geophyllus*, 5.25 U/mL and *Penicillium expansum*, 6.37 U/mL) are mentioned in Table 1.

### Determine optimal conditions for $\alpha$ -amylase production

Describe in sentence-Maximum amount of  $\alpha$ -amylase (8.14 U/mL) by *Aspergillus niger* (Table 2) was produced under conditions: : M1, culture medium; yeast extract , nitrogen source molasses (how much), carbon source; 72 h, incubation time period; 6.5, initial pH of culture medium; 40° C, incubation temperature;  $5 \times 10^6$  conidia, inoculum size; 150 rev/min, agitation rate. Maximum amount of  $\alpha$ -amylase by other fungi (*Aspergillus fumigatu*, 7.01 U/mL, *Penicillium notatum*, 6.58 U/mL; *Penicillium lilacinum*, 7.68 U/mL; optimal conditions for the production of  $\alpha$ -amylase by *Mucor geophyllus*, 4.87 U/mL and *Penicillium expansum*, 5.62 U/mL) are mentioned in Table 2.

Table 2. Optimal conditions for the production of alpha amylase by various fungi

Fungi	Culture medium	Incubation time period	Carbon source	Nitrogen source	Incubation Temperature	Initial pH of Medium	Inoculum Size	Agitation Rate	Optimized $\alpha$ -amylase activity
		h			° C		Conidia/mL	Rev/min	U/mL
<i>A. niger</i>	M1	72	molasses	yeast extract	40	6.5	$5 \times 10^6$	150	8.14
<i>A. fumigatus</i>	M1	72	Sunflower waste	casein	35	5.5	$6 \times 10^6$	150	7.01
<i>P. notatum</i>	M1	48	molasses	corn steep liquor	30	5.5	$5 \times 10^6$	150	6.58
<i>P. lilacinum</i>	M1	96	molasses	yeast extract	40	7.5	$6 \times 10^6$	200	7.68
<i>M. geophyllus</i>	M1	48	molasses	yeast extract	35	6.5	$5 \times 10^6$	150	4.87
<i>P. expansum</i>	M1	72	molasses	yeast extract	35	5.5	$5 \times 10^6$	150	5.62

## DISCUSSION

The production of enzyme depends upon various parameters such as carbon source, nitrogen source, the strain, culture medium, incubation time period, incubation temperature, initial pH of culture medium, agitation rate and inoculums (Ahmed *et al.*, 2015a; 2015b; 2015c, 2016).. Uma *et al.* (2010) reported Fruit peel as the most appropriate carbon source for the production of invertase by *Aspergillus flavus*, Pomegranate peel by Uma *et al.* (2012) and Sugar cane bagasse by Guimaraes *et al.*, (2007). While for  $\alpha$ - amylase production, Pomegranate peel (Singh *et al.*, 2014), Wheat bran (Khan and Yadav, 2011), and starch ((Saleem and Ebrahim, 2014) were reported as the most appropriate carbon source.

Nitrogen source is also important constituent of Culture medium. Yeast extract (Uma *et al.*, 2010), Yeast extract plus peptone (Olusanya and Oliotula, 1994), Peptone (Belcarz *et al.*, 2000) and Corn steep liquor (Chan *et al.*, 1991) have been reported as the most suitable nitrogen source for the maximum production of invertase while for  $\alpha$ -amylase, Beef extract (Singh *et al.*, 2014), Peptone (Khan and Yadav, 2011), Ammonium nitrate (Matthias, 2013) and Peptone and ammonium sulphate (Saleem and Ebrahim, 2014) have been reported.

Incubation time period is also important parameter for enzyme production. In literature 96 h (Uma *et al.*, 2010),

24 h (Zafar and Aslam, 2013) and 48 h (Mizunaga *et al.*, 1981) are reported as the best incubation time period for the production of invertase while for  $\alpha$ -amylase, 144 h (Singh *et al.*, 2014), 48 h (Khan and Yadav, 2011) and 120 h (Matthias, 2013) have been reported.

Initial pH of culture medium is important for quality and quantity of culture medium. In order to obtain maximum quantity of invertase, initial pH of 5.0 (Uma *et al.*, 2010), 4.0 (Uma *et al.*, 2012) and 8.0 (Qureshi *et al.*, 2012) were reported as the best while for  $\alpha$ -amylase, a pH of 6.0 (Singh *et al.*, 2014), 6.2 (Khan and Yadav, 2011), and 4.0 (Matthias, 2013) have been reported.

Growth of microorganism is influenced by incubation temperature. It has been reported that 30° C (Uma *et al.*, 2010) and 45° C (Qureshi *et al.*, 2012) is the most suitable incubation temperature while for  $\alpha$ -amylase production, 35° C (Singh *et al.*, 2014), 28° C (Khan and Yadav, 2011), 30° C (Saleem and Ebrahim, 2014) and 45° C (Matthias, 2013) have been reported.

## CONCLUSION

The highest quantity of invertase (13.05 U/mL) was obtained by *Penicillium lilacinum* under conditions: CM1, culture medium; yeast extract, nitrogen source; date syrup, carbon source; 96 h, incubation time period; 8, initial pH of culture medium; 40° C, incubation temperature;  $6 \times 10^6$  conidia, inoculum size; 200 rev/min, agitation rate. While the highest amount of  $\alpha$ -amylase (8.14 U/mL) was obtained by *Aspergillus niger* under conditions: : M1, culture medium; yeast extract, nitrogen source; molasses, carbon source; 72 h, incubation time period; 6.5, initial pH of culture medium; 40° C, incubation temperature;  $5 \times 10^6$  conidia, inoculum size; 150 rev/min, agitation rate.

**Acknowledgment:** We thank HEC Pakistan for providing research support through its Project No. 21-826/SRGP/R&D/HEC/2016

## REFERENCES

- Ahmed, K., E.E. Valeem and Qamar-ul-Haq (2015a). Biosynthesis, Purification and Characterization of Commercial Enzyme by *Penicillium expansum* Link. *Pak. J. Bot.*, 47: 1521-1526.
- Ahmed, K., S. Munawar and M. A. Khan (2015b). Cultural conditions for maximum alpha-amylase production by *Penicillium notatum* IBGE 03 using shaken flask technique of submerged fermentation. *Pure and Applied Biology*, 4(3): 306-312
- Ahmed, K., S. Munawar and M. A. Khan (2015c). Increased industrial enzyme production by *Penicillium lilacinum* ibge 04 using shaken flask technique of submerged fermentation. *Science International*, 27(3):2133-2137.
- Ahmed, K., T. Ashraf and S. Naz (2016). Cultural conditions for the increased production of industrial enzyme by *Mucoroglyphus oudem* along with partial purification and characterization. *Int. J. Biol. Biotech.*, 13: 193-201.
- Almeida, A.C.S., L.C. Araujo, A.M. Costa, C.A.M. Abreu, M.A.G.A. Lima and M.L.A.P. F.P. Pahl (2005). Sucrose hydrolysis catalyzed by auto-immobilized invertase into intact cells of *Cladosporium cladosporioides*. *E. J. Biotech.*, 8(1): 54-62.
- Belcarz, A., G. Ginalska, J. Lobarzewski, H. Greppin and J. Fiedurek (2000). The optimization of the liquid affinity chromatography conditions of the extracellular invertase isolation from the *Candida utilis* cultures. *Chromatographia*, 51(1): 121-129.
- Burrell, R.G., C.W. Clayton, M.R. Gallegly and V.G. Litty (1966). Factors affecting the antigenicity of the mycelium of three species of *Phytophthora*. *Phytopathol.*, 56: 422-426.
- Chan, E., C. S. Chen, C. S. Gong and L. F. Chen (1991). Production, separation and purification of yeast invertase as a by-product of continuous ethanol fermentation. *Appl. Microbiol. Biotechnol.*, 36(1): 44-47.
- Dworschack, R. G. and L. J. Wickerham (1961). Production of extracellular and total invertase by *Candida utilis*, *Saccharomyces cerevisiae*, and other yeasts. *Appl. Microbiol.*, 9(4): 291-294.
- Guimarães, L. H. S., H. F. Terenzi, M. L. T. M. Polizeli and J.A. Jorge (2007). Production and characterization of a thermostable extracellular  $\beta$ -D-fructosuranosidase produced by *Aspergillus ochraceus* with agroindustrial residues as carbon source. *Enzyme Microb. Technol.*, 42(1): 52-57.
- Hayashida, S. and Y. Teramoto (1986). Production and characteristics of raw starch digesting  $\alpha$ -amylase from protease negative *Aspergillus ficummutant*. *Appl. Environ. Microbiol.*, 52 (5): 1068-1073.
- Khan, J. A. and S. K. Yadav (2011). Production of alpha amylases by *Aspergillus niger* using cheaper substrates employing solid state fermentation. *Int. J. Plant. Anim. and Environ. Sci.* 3(1): 100-108.
- Kulshrestha, S. (2013). Invertase and its applications- a brief review. *J. Pharm. Res.*, 7(1): 792-797.
- Matthias, O. C. (2013). Optimization of  $\alpha$ -amylase and glucoamylase production from three fungal strains isolated from Abakaliki, Ebonyi State. *Eur. J. Exp. Biol.*, 3(4): 26-34.

- Mamma, D., E. Kourtoglou and P. Christakopoulos (2008). Fungal multienzyme production on industrial by-products of the citrus-processing industry. *Bioresour. Technol.*, 99(7): 2373-2383.
- Mizunaga, T., J. S. Ikacz, L. Rodriguez, R. A. Hackel and J. O. Lampen (1981). Temperature-sensitive forms of large and small invertase in a mutant derived from a Suc1 strain of *Saccharomyces cerevisiae*. *Mol. Cell Biol.*, 1(5): 460-468.
- Olusanya, O. and P. O. Olutiola (1994). The purification and characterization of intracellular invertase obtained from pathogenic *Escherichia coli*. *Afr. J. Med. Sci.*, 23(3): 291-299.
- Poonawalla, F.M., K.L. Patel and M.R.S. Iyenger (1965). Invertase Production by *Penicillium chrysogenum* and Other Fungi in Submerged Fermentation. *Appl. Microbiol.*, 13(5): 749-754.
- Qureshi, A. S., I. Khusk, M. A. Bhutto, M. U. Dahot, I. Haq, S. Bano and H. Iqbal (2012). Production and partial characterisation of invertase from *Mucor geophyllus* EFRL 03. *Af. J. Biotech.*, 11(47): 10736-10743.
- Saleem A. and M. K. H. Ebrahim (2014). Production of amylase by fungi isolated from legume seeds collected in Almadinah Almunawwarah, Saudi Arabia. *J. Taibah Uni. Sci.*, 8: 90-97.
- Singh, S., S. Singh, V. Bali, L. Sharma and J. Mangla (2014). Production of fungal amylases using cheap, readily available agri-residues, for potential application in textile industry. *Bio Med Res. Int.*, p.1-9 (<http://dx.doi.org/10.1155/2014/215748>).
- Sundarram A. and T. P. K. Murthy (2014).  $\alpha$ -amylase production and applications: a review. *J. App. & Environ. Microbiol.*, 2(4): 166-175.
- Souza, M.J., C., Alves-Araújo, A., Pacheco, M.J., Almeida, I. Spencer-Martins and C. Leão (2007). Sugar utilization patterns and respiro-fermentative metabolism in the baker's yeast. *Torulaspora delbrueckii*. *Microbiol.*, (153): 898-904.
- Uma, C., D. Gomathi, C. Muthulakshmi and V. K. Gopalakrishnan (2010). Production, purification and characterization of invertase by *Aspergillus flavus* using fruit peel waste as substrate. *Adv. Biol. Res.* 4(1): 31-36.
- Uma, C., D. Gomathi, G. Ravikumar, M. Kalaiselvi and M. Palaniswamy (2012). Production and properties of invertase from a *Cladosporium cladosporioides* in SmF using pomegranate peel waste as substrate. *A. Pac. J. Trop. Biomed.*, 144: 605-611.
- Zafar, F. and A. Aslam (2013). A comparative study of nutritional and environmental factors affecting extracellular and intracellular invertase production in *Candida utilis*. *Pak. J. Bot.*, 45(2): 681-686.

(Accepted for publication February 2018)