

NEOECHINORHYNCHUS RAFIAE SP. N. (ACANTHOCEPHALA: NEOECHINORHYNCHIDAE) FROM MUGIL CEPHALUS LINN., 1758 IN KARACHI COAST, PAKISTAN

Faiza Saleem¹, Aly Khan^{2*}, Nasira Khatoon¹, Samina Waheed¹ and Noor-un-Nisa³

¹Department of Zoology, University of Karachi, Karachi-75270, Pakistan

²CDRI, Pakistan Agricultural Research Council, University of Karachi, Karachi-75270, Pakistan

³Vertebrate Pest Control Institute, Southern-zone Agricultural Research Centre, PARC, Karachi University Campus, Karachi-75270, Pakistan

*Corresponding author's e-mail: aly.khan@hotmail.com

ABSTRACT

Mugil cephalus Linnaeus, 1758 is found in coastal subtropical and tropical waters around the world, being an important variety of fish. It is fished as well as farmed. A new species of thorny-headed worm (Acanthocephala: Neoechinorhynchidae) namely *Neoechinorhynchus rafiae* is being reported. The genus is distributed worldwide and parasitizes turtles and fish which is being reported from Karachi coast, Sindh, Pakistan. The new species *N. rafiae* has unique characteristics of the hooks of first and second row/circle of the same size thus differs from its congeners on this important character. The present species comes close to *N. oreini* Fotedar, 1968 but differs in a number of characters. The prominent feature observed in the present species is the presence of para-receptacle structure (PRS) prominent along ventral side of receptacle.

Keywords: *Neoechinorhynchus rafiae* sp.n., *Mugil cephalus*, Karachi coast, Pakistan.

INTRODUCTION

Mulletts are fish of the family Mugilidae which usually feed on zooplankton and algae. *Mugil cephalus* (Linnaeus, 1758) is found in coastal subtropical and tropical waters (Zahid *et al.*, 2015). It is an important food fish and is fished as well as farmed (FAO, 2016). Its roe is salted, dried and compressed as a speciality food in Italy, Japan, Korea, Turkey and Egypt. Fried Mullet is the most popular form but canned, baked and smoked are also popular in fish food outlets. Mullet are usually filleted and the remaining frames are mostly used for fish stock in stews and chowders (Anonymous, 2007). *Mugil cephalus* is a popular food item in the province of Sindh. Therefore, the interest was developed to investigate this fish for the presence of Acanthocephala parasites.

MATERIALS AND METHODS

Fish (*Mugil cephalus* Linnaeus, 1758) were caught alive in January 2019 and brought to Parasitology Laboratory, Department of Zoology, University of Karachi, Pakistan. The fish were dissected and examined for collection of Acanthocephala parasites. During examination five parasites were recorded from a single fish small intestine. These specimens were fixed in cold 70% ethanol so that the proboscis come out of the body. Later these specimens were gently pressed between two slides for 24 hours and fixed in F.A.A solution, stained with Mayer's carmalum, dehydrated in graded series of alcohol, cleared in clove oil and rinsed in xylene. Finally, the specimens were permanently mounted in Canada balsam for further study, drawing were made with the aid of camera Lucida. Measurements are in millimeters (mm) and length by with. Photomicrographs were prepared using Nikon photomicroscope. Holotype and Paratype slides are in collection of senior author (F.S).

RESULTS

***Neoechinorhynchus rafiae* sp.n.**

(Figs. 1-3)

Host: *Mugil cephalus* Linnaeus, 1758
Location: Small intestine
Locality: Karachi coast (24°51' 39.4776N and 66°59' 25.8036"E) Pakistan
No. of specimens recovered: 8 from 3 hosts
No. of hosts examined: 10




Fig. 1. *Neoechinorhynchus raftiae* sp.n., **a.** Entire specimen, holotype; **b.** Anterior end; **c.** Hooks enlarged; **d.** Eggs; **e.** Posterior end of female.

DESCRIPTION

Neoechinorhynchidae, Neoechinorhynchinae with characters of the genus. Trunk medium size with sexual dimorphism in common characters. Widest in testicular region in males and in females just behind proboscis receptacle. Body medium size and cylindrical. Male posterior end slightly curved. Shared characters slightly larger in females as compared to males. Giant nuclei not marked. Proboscis small in size as compared to trunk. Proboscis hooks in all the three circles with roots. Proboscis hooks of first and second circle almost same in size. Neck short. Proboscis receptacle elongated with single wall with triangular cephalic ganglion at its base. Para-receptacle structure seen along ventral side of proboscis receptacle both in male and female specimens. Eggs without polar prolongation of the fertilization membrane. Lemnisci slightly subequal and elongated.




Fig. 2: Photomicrograph showing rows of hooks (x 400).

Fig. 3: Photomicrograph showing anterior region of the thorny-headed worm (x 100).

Male (based on six mature specimens). Body size 4.0-4.6 by 0.22-0.27. Proboscis small in size measuring 0.24-0.27 by 0.20-0.22. Hooks of first row measuring 0.013-0.015 by 0.0017-0.0019; second row 0.013-0.015 by 0.0017-0.0018 and third row 0.0076-0.0078 by 0.0014-0.0017. Neck 0.072-0.015 by 0.18-0.23. Proboscis receptacle 0.30-0.42 by 0.18-0.22. Lemnisci left 1.25-1.29 by 0.08-0.11. Lemnisci right 1.25-1.28 by 0.08-0.11. Testis anterior 0.31-0.38 by 0.14-0.18. Testis posterior 0.31-0.38 by 0.14-0.18. Cement gland elongated measuring 0.38-0.42 by 0.07-0.08. Saeftigen's pouch measuring 0.36-0.41 by 0.07-0.08. Cement reservoir elongated measuring 0.75-0.78 by 0.09-0.12. Bursa eversible measuring 0.25-0.28 by 0.16-0.19.

Females (based on two mature specimens). Body size 5.4-5.8 by 0.47-0.59. Proboscis 0.25-0.27 by 0.22-0.23. Hooks of the first row measuring 0.014-0.016 by 0.0017-0.0019; hooks of second row measuring 0.014-0.015 by 0.0017-0.0019 and the third row measuring 0.0077-0.0078 by 0.0015-0.0018. Neck 0.079-0.16 by 0.18-0.23. Proboscis receptacle 0.30-0.42 by 0.21-0.23. Lemnisci left 1.26-1.28 by 0.08-0.11. Lemnisci right 1.26-1.28 by 0.08-0.11. Uterus length measuring 0.90-0.95; vaginal funnel width 0.40-0.42; vulval sphincter 0.30-0.32; vaginal bulb measuring 0.36-0.39 by 0.10-0.14. Sub-terminal gonospore. Mature eggs ovoid and no polar prolongation 0.30-0.34 by 0.15-0.18.

DISCUSSION

Neoechinorhynchid life cycle in Mulletts include ostracods (seed shrimp) as intermediate hosts along with crayfish and snails presumably acting as paratenic host (Merritt and Prat, 1964). Cable and Dill (1967) studied that life cycles for those neoechinorhynchids infecting mullet after considering the constituency in Acanthocephalan life cycle, probably acquire helminths from feeding on snails, ostracods and copepods.

The most striking feature in the present specimens is that in both sexes the size of first and second circle hooks are same in size. While in other species are unequal namely *N. aminulhaquei* Chandra, 1983; *N. argentatus* Chandra

et al., 1987; *N. bangoni* Tripathi, 1959; *N. chilkaensis* Podder, 1937; *N. brayi* Bilqees *et al.*, 2011; *N. cirrhinae* Gupta and Jain, 1979; *N. dattai* Golvan, 1994; *N. devdevi* (Datta, 1936) Kaw, 1951; *N. elongatus* Tripathi, 1959; *N. glyptosternumi* Fotedar and Dhar, 1977; *N. formosanus* (Harada, 1938) Kaw, 1951; *N. gibsoni* Khan and Bilqees, 1989; *N. hutchisoni* Datta, 1936; *N. johnii* Yamaguti, 1939; *N. kallarensis* George and Nadakal, 1978; *N. karachiensis* Bilqees, 1972; *N. macrorchis* Shaikh *et al.*, 2011; *N. nawazi* Naqvi *et al.*, 2012; *N. nickoli* Khan *et al.*, 1999; *N. nematalosi* Tripathi, 1959; *N. nickoli* Khan *et al.*, 1999; *N. ovalis* Tripathi, 1959; *N. rigidus* (Van Cleave, 1928) Kaw, 1951; *N. roonwali* Datta and Soota, 1963; *N. sootai* Bhattacharya, 1999; *N. topseyi* Podder, 1937 while in *N. cyanophlyctis* Kaw, 1951 they are all of equal length 0.014-0.025 long.

The present specimens comes close to *N. oreini* Fotedar, 1968 reported from *Oreinus sinuatus* from Arapat stream, Anant Nag Jammu and Kashmir, India in having equal size of first and second row of hooks but *N. oreini* differs in having larger hooks size (0.078-0.088 by 0.073-0.085). Body size (8.0-11.75 by 0.9-1.45 male and female 11.5-16.8 by 1.2-1.7) is also larger as compared to the present species. Proboscis in present specimens is larger as compared to *N. oreini* (0.13-0.18 by 0.10-0.16). Lemnisci in present specimens are smaller as compared to *N. oreini* (1.59-2.9 by 1.32-2.68). In *N. oreini* the difference in the testes anterior testis size (0.80-1.5 by 0.56-0.70) and posterior testis (0.69-1.4 by 0.49-0.6) have a lot of difference while in the present specimens they are almost of the same size anterior testis (0.31-0.38 by 0.14-0.18) and posterior testis (0.31-0.38 by 0.14-0.18).

In body size the present specimens are larger as compared to *N. aminulhaquei* Chandra, 1983; *N. cirrhinae* Gupta and Jain, 1979; *N. cyanophlyctis* Kaw, 1951; *N. ovalis* Fotedar, 1968; *N. sootai* Bhattacharya, 1999; *N. nematalosi* Tripathi, 1959; *N. ovale* Tripathi, 1959; and *N. zacconis* Yamaguti, 1935.

The proboscis in present specimens is smaller as compared to *N. idahoensis* Amin and Heckmann, 1992; *N. dimorphospinus* Amin and Sey, 1996; *N. pimelodi* Brasil-Sato and Pavanelli, 1998; *N. didelphis* Amin, 2001; *N. iraqensis* Amin *et al.*, 2001; *N. qatarensis* Amin *et al.*, 2002; *N. zabensis* Amin *et al.*, 2003; *N. buckneri* Amin and Heckmann, 1992; *N. longinucleatus* Amin *et al.*, 2011; *N. colastinensis* Arredondo and Gil de Pertierra, 2012; *N. vittiformis* Smales, 2013; *N. bryanti* Smales, 2013; *N. veropesoi* Melo *et al.*, 2013 and *N. yamagutii* Tkach *et al.*, 2014.

Giant-Hypodermic nuclei are present in *N. daleri* Khan *et al.*, 2020; *N. rigidis* (Van Cleave, 1928) Kaw, 1951; *N. roonwali* Datta and Soota, 1963; *N. devdevi* (Datta, 1936) Kaw, 1951; *N. chilkaensis* Podder, 1937; *N. oreini* Fotedar, 1968; *N. dattai* Golvan, 1994; *N. formosanus* (Harada, 1938) Kaw, 1951; *N. tylosuri* Yamaguti, 1939; *N. inermis* Brito-Porto *et al.*, 2017; *N. personatus* Tkach *et al.*, 2014; *N. ponticus* Amin *et al.*, 2020; *N. daleri* Khan *et al.*, 2020 and *N. qatarensis* Amin *et al.*, 2002 while were not observed in present specimens.

At higher magnification the para-receptacle structure (PRS) was observed which was first described by Amin *et al.* (2002) in *Neoechinorhynchus qatarensis* from the marine blue-barred flame parrot fish, *Scarus globban* Forskal, 1775 in Qatari waters of Arabian gulf. The para-receptacle structure (PRS) prominent along ventral side of receptacle, connecting anteriorly to body wall as earlier described by Amin *et al.* (2007) later reported that PRS as an oblong capsule enclosing a column of nucleated core (NC) cells appeared lighter coloured as compared to darker walls of PRC. It is believed that in North American continent has the largest number of species known in this genus (Amin, 2002).

Due to morphological and morphometric differences between the present specimens and earlier described species of the genus *Neoechinorhynchus* Stiles and Hassall, 1905 a new species *N. rafiae* is proposed. Name of the new species is in honour of Dr. Rafia Rehana Ghazi a distinguished Parasitologist of Pakistan.

REFERENCES

- Amin, O.M. (2001). *Neoechinorhynchus didelphis* sp.n. (Acanthocephala: Neoechinorhynchidae) from the Redfin Pickerel, *Esox americanus*, in Georgia, U.S.A. *Comp. Parasitol.*, 68: 103-107.
- Amin, O.M. (2002). Revision of *Neoechinorhynchus* Stiles and Hassall, 1905 (Acanthocephala: Neoechinorhynchidae) with keys to 88 species in two subgenera. *Syst. Parasitol.*, 53: 1-18.
- Amin, O.M. and O. Sey (1996). Acanthocephala from Arabian Gulf fishes off Kuwait, with descriptions of *Neoechinorhynchus dimorphospinus* sp. n. (Neoechinorhynchidae), *Tegorhynchus holospinus* sp. n. (Illiosentidae), *Micraacanthorhynchina kuwaitensis* sp. n. (Rhabdinorhynchidae), and *Slendrorhynchus breviclaviproboscis* gen. n., sp. n. (Diplosentidae); and key to species of the genus *Micraacanthorhynchina*. *J. Helminthol. Soc. Wash.*, 63: 201-210.
- Amin, O.M. and R.A. Heckmann (1992). Description and pathology of *Neoechinorhynchus idahoensis* n. sp. (Acanthocephala: Neoechinorhynchidae) in *Catostomus columbianus* from Idaho. *J. Parasitol.*, 78: 34-39.

- Amin, O.M., Ha, N.V. and Ha, D.N. (2011). First report of *Neoechinorhynchus* (Acanthocephala: Neoechinorhynchidae) from marine fish of the eastern seaboard of Vietnam, with the description of six new species. *Parasite*, 18: 21-34.
- Amin, O.M., M. Sharifdini, R.A. Heckmann, N. Rubtsova and H.J. Chine (2020). On the *Neoechinorhynchus agilis* (Acanthocephala: Neoechinorhynchidae) complex, with a description of *Neoechinorhynchus ponticus* n. sp. from *Chelon auratus* in the Black Sea. *Parasite*, 27: 19-36.
- Amin, O.M., M.F.A. Saoud and K.S.R. Alkuwari (2002). *Neoechinorhynchus qatarensis* sp. n. (Acanthocephala: Neoechinorhynchidae) from the Blue-barred flame parrot fish, *Scarus gobban* Forsskal, 1775, in Qatar waters of the Arabian Gulf. *Parasitol. Int.*, 51: 171-196.
- Amin, O.M., R.A. Heckmann and M.D. Standing (2007). Structural-functional relationship of the para-receptacle structure in Acanthocephala. *Comp. Parasitol.*, 74: 383-387.
- Amin, O.M., R.S.S. Al-Sady, F.T. Mhaisen and S.F. Bassat (2001). *Neoechinorhynchus iraqensis* sp. n. (Acanthocephala: Neoechinorhynchidae) from the freshwater Mullet, *Liza abu* (Heckel), in Iraq. *Comp. Parasitol.*, 68: 108-111.
- Amin, O.M., S.M.A. Abdullah and F.T. Mhaisen (2003). *Neoechinorhynchus* (*Neoechinorhynchus zabensis* sp. n. (Acanthocephala: Neoechinorhynchidae) from freshwater fish in northern Iraq. *Folia Parasitol.*, 50: 293-297.
- Anonymous. (2007). Istituto Veneto di scienze, Lettere ed arti environmental database on the pool of Venice. Archived from the original in 2010-10-13. Retrieved 2007-05-23.
- Arredondo, N.J. and A.A. Gil de Pertierra (2012). A new species of *Neoechinorhynchus* (Eoacanthocephala: Neoechinorhynchidae) from *Pachyurus bonariensis* (Perciformes: Sciaenidae) from the Parana River basin in Argentina, with comments on two other species of the genus. *Revue Suisse De Zoologie*, 119: 425-439.
- Bhattacharya, S.B. (1999). State Fauna Series, 4. Fauna of Meghalaya (Part 9): Acanthocephala. Zoological Survey of India, Calcutta. pp. 359-392.
- Bilqees, F.M. (1972). Description of two acanthocephala including a new species *Neoechinorhynchus karachiensis* (Neoechinorhynchidae: Neoechinorhynchida) from marine fishes of Karachi. *Sind. Uni. Res. J.*, 6: 93-100.
- Bilqees, F.M., G.S. Shaikh and A. Khan (2011). *Neoechinorhynchus brayi* sp.n. (Acanthocephala: Neoechinorhynchidae) in freshwater fish *Catla catla* from Sindh, Pakistan. *Int. J. Biol. Biotech.*, 8: 491-494.
- Brasil-Sato, M.de.C. and G.C. Pavanelli (1998). *Neoechinorhynchus pimelodi* sp. n. (Eoacanthocephala, Neoechinorhynchidae) parasitizing *Pimelodus maculatus* Lacepede, "Mandi Amarelo" (Siluroidei, Pimelodidae) from the Basin of the Sao Francisco River, Tres Marias, Minas Gerais, Brazil. *Revta Bras. Zool.*, 15: 1003-1011.
- Brito-Porto, D., A.K. Silva-de Souza and J.C. de Oliveira-Malta (2017). A new species of *Neoechinorhynchus* (Eoacanthocephala: Neoechinorhynchidae) from the freshwater fish *Ageneiosus inermis* (Siluriformes) in the Brazilian Amazon. *Rev. Mex. Biodivers.*, 88: 798-800.
- Cable, R.M. and W.T. Dill (1967). The morphology and life history of *Paulisentis fractas* Van Cleave and Bangham, 1949 (Acanthocephala: Neoechinorhynchidae) *J. Parasitol.*, 53: 810-817.
- Chandra, K.J. (1983). On *Neoechinorhynchus aminulhaquei* n. sp. (Acanthocephala) from a freshwater fish, *Mystus vittatus* (Bloch). *Indian J. Zool.*, 24: 131-135.
- Chandra, K.J., K.H. Rao and K. Shyamsundari (1987). On *Neoechinorhynchus argentatus* new species, an Acanthocephalan parasite from a marine fish of Waltair. *Revta. Iber. Parasit.*, 45: 49-52.
- Datta, M.N. (1936). Scientific results of the Yale North India Expedition Biological Report no.20. Helminth parasites of fishes from North India, with special reference to acanthocephalans. *Rec. Ind. Mus.*, 38: 211-229.
- Datta, M.N. and T.D. Soota (1963). A new species of Acanthocephala *Neoechinorhynchus roonwali* from Afghanistan, and notes on *Pomphorhynchus kashmirensis* Kaw. *Rec. Indian Mus.*, 59: 35-39.
- FAO. (2016). Species fact sheets-*Mugil cephalus* (Linnaeus, 1758) fao.org. Archived from the original on 2015-11-05. Retrieved 2016-04-30.
- Forsskal, P. (1775). Descriptions animalium avium, amphibiorum, piscium, insectorum, vermium, quae in itinere orientali observavit Petrus Forsskal. Post mortem auctoris edidit Carsten Niebuhr. Adjuncta est material medica Kahirina atque tabula maris rubri geographica. pp. 1-164.
- Fotedar, D.N. (1968). New species of *Neoechinorhynchus* Hamann, 1882, from *Oreinus sinuatus*, fresh water fish in Kashmir. *Kash. Sci.*, 5: 147-152.
- Fotedar, D.N. and R.L. Dhar (1977). Wide range infections of *Neoechinorhynchus* Hamann, 1892 in freshwater fishes of Jammu and Kashmir. In all India Symp. on Helminthology, Srinagar 8-11 Aug., 1977. pp. 36-37.
- George, P.V. and A.M. Nadakal (1978). Two new species of Acanthocephala from marine fishes of Kerala. *Aqua. Biol.*, 3: 91-97.
- Golvan, Y.J. (1994). Nomenclature of the Acanthocephala. *Res. Rev. Parasit.*, 54: 135-205.

- Gupta, N.K. and M. Jain (1979). On *Neoechinorhynchus cirrhinae* sp.n., an acanthocephalan parasite from a freshwater fish at Chandigarh. *Helminthologia*, 16: 147-151.
- Harada, I. (1938). Acanthocephala of Formosa. *Annot. Zool. Jap.*, 17: 419-427.
- Kaw, B.I. (1951). Studies in helminthology: Helminth parasites of Kashmir. Part II. Acanthocephala. *Ind. J. Helm.*, 3: 117-132.
- Khan, A. and F.M. Bilqees (1989). On a new acanthocephalan *Neoechinorhynchus gibsoni*, new species from a freshwater fish, *Labeo rohita* (Ham.). *Proc. Pakistan Congr. Zool.*, 9: 259-264.
- Khan, A., F. Saleem, N. Khatoon, S. Waheed and S.N. Das (2020). *Neoechinorhynchus daleri* n. sp. (Acanthocephala: Neoechinorhynchidae) in the marine fish *Mugil cephalus* (Linnaeus, 1758) from Karachi coast, Pakistan. *Fuuast J. Biol.*, 10: 1-7.
- Khan, A., F.M. Bilqees, Noor-Un-Nisa, R.R. Ghazi and Ata-Ur-Rahim (1999). *Neoechinorhynchus nickoli*, new species (Acanthocephala: Neoechinorhynchidae) from *Labeo boga* (Ham.) of Punjab, Pakistan. *Pakistan J. Zool.*, 31: 241-243.
- Linnaeus, C. (1758). *Systema naturae per regna trianaturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decimal, reformata. Holmiae. (Salvius). Tomus I: 1-824.*
- Melo, F.T.V., P.A.F.B. Costa, E.G. Giese, S.L. Gardner and J.N. Santos (2013). A description of *Neoechinorhynchus (Neoechinorhynchus) veropesoi* n. sp. (Acanthocephala: Neoechinorhynchidae) from the intestine of the silver croaker fish *Plagioscion squamosissimus* (Heckel, 1840) (Osteichthyes: Sciaenidae) off the east coast of Brazil. Faculty publications from the Harold W. Manter Laboratory of Parasitology. pp. 763.
- Merritt, S.V. and I. Pratt (1964). The life history of *Neoechinorhynchus rutili* and its development in the intermediate host (Acanthocephala: Neoechinorhynchidae). *J. Parasitol.*, 50: 394-400.
- Naqvi, S.M.H.M., A. Khan, R.R. Ghazi and Noor-un-Nisa (2012). A new Acanthocephala, *Neoechinorhynchus nawazi* sp.nov. (Neoechinorhynchidae), from a freshwater fish (*Cirrhinus mrigla* Ham.). *Int. J. Biol. Biotech.*, 9: 195-199.
- Podder, T.N. (1937). A new species of Acanthocephala; *Neoechinorhynchus topseyi* n. sp., from a Calcutta fish, *Polynemus heptadactylus* (Cuv.et val). *Parasit.*, 29: 365-369.
- Shaikh, G.S., A. Khan and F.M. Bilqees (2011). On a new Acanthocephala *Neoechinorhynchus macrorchis* sp.nov. (Neoechinorhynchidae) from a freshwater fish of Haleji lake Sindh, Pakistan. *Int. J. Biol. Biotech.*, 8: 333-336.
- Smales, L.R. (2013). A review of the genus *Neoechinorhynchus* (Acanthocephala: Neoechinorhynchidae) from Australia with the description of two new species. *J. Parasitol.*, 99: 1106-1112.
- Stiles, C.W. and A. Hassall (1905). The determination of generic types and a list of roundworm genera, with their original and type species. *Bulletin of the Bureau of Animal Industry, United States Department of Agriculture*, 79: 1-150.
- Tkach, Ie. V., V.L. Sarabeev and L.S. Shvetsova (2014). Taxonomic status of *Neoechinorhynchus agilis* (Acanthocephala, Neoechinorhynchidae), with a description of two new species of the genus from the Atlantic and Pacific Mulletts (Teleostei, Mugilidae). *Vestn. Zool.*, 48: 291-306.
- Tripathi, Y.R. (1959). Studies on parasites of Indian fishes, V. Acanthocephala. *Rec. Ind. Mus.*, 54: 61-99.
- Van Cleave, H.J. (1928). Two new genera and species of Acanthocephala from fishes of India. *Rec. Indian Mus.*, 30(2): 147-149.
- Yamaguti, S. (1935). Studies on the helminth fauna of Japan Part 8 Acanthocephala, I. *Jap. J. Zool.*, 6: 247-278.
- Yamaguti, S. (1939). Studies on the helminth fauna of Japan. Part 29th. Acanthocephala II. Japan. *Jap. J. Zool.*, 8: 317-351.
- Zahid, H., N. Bano, Z. Masood, Musarrat-Ul-Ain, R.Y. Farooq and W. Razaq (2015). Scale surface structure of *Mugil cephalus* (Teleostei; Mugilidae) using scanning electron microscopy (SEM). *Biological Forum – An International Journal*, 7:1845-1848.

(Accepted for publication March 2021)