CLIMATE CHANGE AND DATE PALM CULTIVATION: CHALLENGES AND ADAPTATIONS IN PAKISTAN

Tahira Jatt

Department of Botany, Shah Abdul Latif University, Khairpur, Pakistan
Corresponding Author: tahira@salu.edu.pk

Received: September 10, 2025; Accepted: October 02, 2025

ABSTRACT

The date palm (*Phoenix dactylifera* L.) is among the oldest cultivated fruit trees and holds immense nutritional, cultural, and economic importance in arid and semi-arid regions. Pakistan is among the world's leading producers of date palm, with major orchards concentrated in Sindh, Balochistan, Southern Punjab and parts of Khyber Pakhtunkhwa. Intensifying climate risks heat waves, erratic and extreme monsoon rainfall, drought, salinity, and flood disasters are reshaping production risks along the entire value chain from flowering to postharvest drying. This review synthesizes recent evidence on observed and projected climate trends relevant to date palm agro-ecosystems in Pakistan and examines their impacts on phenology, fruit set and quality, pests and diseases, water use, postharvest processing, and market access. The paper then outlines adaptation strategies climate-smart irrigation, salinity and heat management, cultivar choice and breeding, tissue culture and orchard rehabilitation, climate-informed integrated pest management, and resilient postharvest systems supported by policy and value-chain interventions. Collectively, these measures can reduce climate risk and stabilize yields and quality, provided they are embedded within local institutions and backed by targeted investments, extension, and climate information services.

Key words: date palm, climate change, impact, management.

INTRODUCTION

The date palm (*Phoenix dactylifera* L.) is one of the most ancient fruit-bearing trees known to humanity, with deep cultural roots and historical significance across the Middle East, North Africa, and South Asia. Often referred to as the "Tree of Life" and the "Blessed Tree," it is frequently mentioned in historical texts, including more than 20 references in the Holy Qur'an. Due to its resilience and productivity under harsh environmental conditions, date palm serves as a staple food, a source of income, and an integral part of dryland agro-ecosystems (Jatt *et al*., 2019).

Date palm has deep agro-ecological, cultural, and economic roots in Pakistan. The country consistently ranks among the top global producers, and districts such as Khairpur (Sindh), Kech/Panjgur (Balochistan), Dera Ismail Khan (Khyber Pakhtunkhwa) and Dera Ghazi Khan (Punjab) are major hubs of both fresh and dried date trade. Dates contribute livelihoods to smallholders and laborers and supply domestic markets as well as exports in semi-processed forms (chuhara) and increasingly as premium table dates. In Khairpur alone, associated residues represent a sizable bioresource stream, underscoring the crop's broader bioeconomy potential (Abul-Soad, 2010; Solangi *et al*., 2014; TDAP, 2025a).

Yet the biophysical envelope that sustained traditional production is shifting. Pakistan has experienced a rise in mean temperatures and more frequent extremes; 2023's national mean temperature was ~0.5 °C above average, while recent years brought record-breaking floods and severe heat waves that disrupted agriculture (PMD, 2024; You *et al*., 2024: Maitlo *et al*., 2024a). The 2022 floods devastated Sindh home to Pakistan's largest date orchards submerging cropland and ravaging rural infrastructure

1. Global and Regional Distribution

Globally, over 5000 cultivars of date palm are being cultivated. In Pakistan, the crop is grown across all provinces, with Sindh, Balochistan, and Punjab contributing significantly to national production. Khairpur district in Sindh is considered the biodiversity hub for date palms, housing over 400 unique cultivars. However, many of these varieties are highly sensitive to monsoon rains, which often coincide with the fruit ripening season, leading to substantial postharvest losses (Jatt *et al*., 2023; Abul-Soad *et al*., 2015).

2. Nutritional and Economic Importance

Date fruits are rich in carbohydrates (primarily glucose and fructose), dietary fiber, essential minerals (potassium, magnesium, calcium, and iron), and vitamins (particularly B-complex), proteins/aminoacids (fig1). This makes them an excellent source of energy and nutrition. Economically, date palm provides livelihoods to millions of people across Pakistan and plays a key role in food security, particularly in arid rural regions. (Jatt *et al*, 2025; Bentrad *et al*., 2020).

3. Value-Added Products of Date Palm

The versatility of date fruit is reflected in the wide range of products derived from it, including: Dehydrated dates, Date dry powder, Date paste and jam, Date chutney, Date syrup and juice, Date biscuits, halva, and preserves, Date ice cream and delight, Development of these products not only enhances the shelf life of the fruit but also contributes to value chain diversification and export potential. Additionally date palm trunk is used as pillar and roof in villages homes, leaves are used for cover of the roof of houses, mats hand fans, are made from leaves which widely used in homes (Jatt *et al*, 2025).

4. Genetic and Varietal Diversity in Pakistan

Pakistan's date palm gene pool consists of over 400 cultivars, many of which are indigenous and not yet scientifically characterized. These include elite commercial varieties like Aseel, Begum Jangi, Dhakki, Fasli, and Karbalain. However, there is an urgent need to introduce globally renowned and climate-resilient varieties into the Pakistani ecosystem to counter rising environmental stress and reduce susceptibility to monsoon rains (Jatt *et al*, 2025).

5. Impacts of Climate Change on Date Palm Cultivation

The climate is affecting date palm in Pakistan in multiple ways in terms of their productivity, yield, diseases etc (Table 1). Some major impacts are discussed under:

5.1 Rising Temperatures and Heatwaves

Rising temperatures and the increasing frequency of heatwaves pose significant risks to date palm cultivation in Pakistan. Elevated heat accelerates the fruit ripening process, leading to shortened developmental stages. While this hastens harvest, it often reduces fruit size and sugar accumulation, directly lowering fruit quality. Heat stress during flowering and fruit filling stages particularly compromises pollination success and the final yield. Such climatic changes threaten the economic value of the date crop, which depends heavily on its nutritional quality and marketable appearance. Pakistan has experienced a 0.5°C rise in temperature over the last century.A projected 2°C increase in temperature may reduce date palm productivity by 15–20%. Rising temperatures lead to increased evapotranspiration, necessitating 15–20% more irrigation (Abul-Soad, 2010; IPCC, 2022).

5.2 Erratic Monsoon Rains

Unpredictable and intense monsoon rains have become a serious challenge for date palm growers. Rainfall during ripening stages results in fruit cracking, fermentation, and fungal infections, all of which drastically reduce marketable yield. For instance, the 2022 floods destroyed a major portion of Sindh's Aseel date crop, highlighting the vulnerability of this sector to untimely precipitation (You *et al*., 2024).

5.3 Drought and Water Scarcity

Frequent droughts and declining water availability are critical concerns in Balochistan and southern Punjab, where date palm cultivation is extensive. Drought stress reduces pollination, fruit set, and overall productivity while simultaneously increasing irrigation demands. Limited access to freshwater resources often forces reliance on groundwater, further stressing the sustainability of orchards in these arid zones. Each tree requires around 12,000 litters of water annually; however, due to climate change, water availability has declined by approximately 20% (Ghazzawy *et al*., 2022; PMD, 2024).

5.4 Soil and Water Salinity

Soil and irrigation water salinity are widespread challenges in Sindh and Punjab. Brackish groundwater used for irrigation contributes to salt accumulation in soils, leading to stunted plant growth, leaf drying, and reduced productivity. Salinity not only hampers photosynthesis and nutrient uptake but also accelerates orchard decline if not

managed properly. This represents one of the most persistent climate-aggravated challenges in Pakistan's date-producing belts (FAO, 2022; TDAP, 2025b).

5.5 Flooding and Waterlogging

Extreme rainfall and flood events submerge orchards, causing waterlogging that damages roots, encourages fungal pathogens, and in severe cases, kills entire plantations. The devastating 2022 floods in Sindh submerged large areas of date orchards, destroying both current harvests and future productivity as tree mortality increased. Such events disrupt the continuity of cultivation cycles and pose long-term risks for growers (FAO, 2022; Khan, 2024).

5.6 Increased Pests and Diseases

Climate change also influences pest and disease dynamics. Warmer climates favor the proliferation and spread of pests such as the red palm weevil (*Rhynchophorus ferrugineus*), a destructive pest responsible for severe yield losses. Outbreaks of red palm weevil have been increasingly reported in Sindh, threatening orchard sustainability and requiring integrated pest management measures. Changing climatic conditions are expected to worsen this trend, making pest control more challenging (Soomro *et al*., 2022; Naveed *et al*., 2023).

Table 1. Impact of Climate Change on Date Palm Cultivation in Pakistan.

Factor	Impact on Date Palm	Evidence/Notes	Reference
Rising Temperatures / Heatwaves	Faster fruit ripening; shortened developmental stages; may reduce fruit size and sugar accumulation	Heat stress accelerates maturity but lowers fruit quality	Abul-Soad (2010); IPCC (2022)
Erratic Monsoon Rains	Fruit cracking, fermentation, fungal growth, and major yield loss during ripening	2022 floods destroyed large share of Sindh's Aseel crop	Dawn (2022); You *et al*., (2024)
Drought and Water Scarcity	Reduces pollination, fruit set, and overall yield; increases irrigation demand	Critical in Balochistan and southern Punjab	Ghazzawy *et al*. (2022); PMD (2024)
Soil and Water Salinity	Stunted growth, leaf drying, reduced productivity	Salinity from brackish groundwater is widespread in Sindh and Punjab	FAO (2022); TDAP (2025b)
Flooding / Waterlogging	Orchard submergence causes tree death and root rot; harvest disruption	2022 Sindh floods submerged large orchards	FAO (2022); Khan (2024)
Increased Pests (e.g., Red Palm Weevil)	Higher infestation and spread under warmer climate; severe yield loss	Red palm weevil outbreaks reported in Sindh	Soomro *et al*. (2022); Naveed *et al*. (2023)
Postharvest Losses	Traditional sun-drying disrupted by untimely rains; increases fungal contamination	Urgent need for solar dryers and cold storage	Solangi *et al*. (2014); TDAP (2025b)

5.7 Postharvest Losses

Postharvest management of dates is highly sensitive to weather conditions. Traditional sun-drying methods, widely practiced by farmers, are increasingly disrupted by untimely rains, leading to fungal contamination and spoilage. This significantly reduces storability and export quality. To address this challenge, there is an urgent need to adopt modern techniques such as solar dryers and cold storage facilities, which can safeguard the quality and shelf life of harvested dates in a changing climate (Solangi *et al*., 2014; TDAP, 2025a).

5.8 Yield and Export Losses

A 20–30% reduction in yield is expected by 2050 under current climate change trajectories. Warmer climates have intensified pest and disease outbreaks, increasing their pressure by up to 40%. Climate-induced stress has already caused a 10% reduction in export volumes, affecting the national economy (Jatt *et al* 2025).

5.9 Loss of Genetic Diversity

Climate change, along with unregulated cross-pollination and urban encroachment, is reducing the genetic diversity of native cultivars. This makes the crop more vulnerable to environmental fluctuations and biological stresses (TDAP, 2025b).

5.10 Physiological and Agronomic Impacts

Date palms are highly sensitive during reproductive stages. Heat stress reduces flowering efficiency and pollination success, leading to lower yields. High humidity and unseasonal rainfall during ripening promote fungal infections and poor fruit quality. Salinity stress disrupts osmotic balance, reducing nutrient uptake, stunting growth, and lowering photosynthetic efficiency. Waterlogging from floods suffocates roots and weakens trees, making them vulnerable to diseases. Additionally, climate change alters pest and pathogen dynamics, increasing threats to orchard health (TDAP, 2025b; Maitlo *et al*., 2025).

5.11 Socio-Economic Consequences

Date palm farming provides income to thousands of rural households, especially in Sindh and Balochistan. However, climate-related yield losses reduce farmer earnings, limit export opportunities, and increase post-harvest costs. Pakistan exports dates to the Middle East, Europe, and South Asia, but reduced quality undermines competitiveness in international markets. Smallholder farmers are particularly vulnerable, as they lack access to modern irrigation technologies, crop insurance, or capital to invest in climate-smart practices (TDAP, 2025b).

6. Strategies for Climate Resilience and Sustainable Cultivation

6.1 Irrigation Management

Efficient irrigation is one of the most important strategies for enhancing climate resilience in date palm cultivation. Traditional flood irrigation contributes to water loss, salinity buildup, and inefficient resource use. To address these challenges, the adoption of drip and sprinkler irrigation systems has been recommended, as these methods significantly reduce water wastage while improving yield and fruit quality. Rainwater harvesting techniques can supplement irrigation in arid regions, particularly in Balochistan and Sindh where water scarcity is acute. Moreover, irrigation scheduling based on evapotranspiration data allows farmers to optimize water application according to plant requirements and prevailing climatic conditions, reducing stress on orchards (FAO, 2022).

6.2 Soil Conservation

Soil conservation practices are critical to sustaining the productivity of date palm plantations under changing climatic conditions. Conservation tillage, which minimizes soil disturbance, helps preserve soil organic matter and improves water retention. Organic mulching reduces surface evaporation, regulates soil temperature, and adds nutrients to the soil as it decomposes. Additionally, cover cropping between date palm rows enhances soil structure, prevents erosion, and contributes to biodiversity. These practices improve overall soil health and provide resilience against climate-induced stresses such as drought and salinity (Maitlo *et al.,* 2024b; Hussain *et al*., 2021).

6.3 Integrated Pest and Disease Management (IPM)

Climate change has intensified pest and disease challenges in date palm orchards, particularly outbreaks of the red palm weevil and fungal pathogens. Integrated pest management (IPM) offers a sustainable solution by combining biological control agents, resistant cultivars, and minimal use of chemical pesticides. Biological controls, such as entomopathogenic fungi and pheromone traps, reduce pest populations while minimizing environmental impacts. Resistant cultivars provide long-term protection, and judicious chemical use prevents pest resistance. By integrating these approaches, IPM reduces yield losses and maintains ecological balance in orchards (Soomro *et al*., 2022; Naveed *et al*., 2023).

6.4 Breeding for Climate-Resilient Varieties

The development of climate-resilient date palm varieties is essential to sustain production in the face of rising temperatures, drought, and salinity stress. Both conventional breeding and modern molecular techniques, such as marker-assisted selection, can be used to identify and propagate cultivars with enhanced tolerance to abiotic and biotic stresses. Efforts to breed drought- and disease-resistant varieties can improve orchard resilience and farmer profitability. Such breeding programs also contribute to reducing reliance on chemical inputs, thereby supporting environmentally sustainable cultivation (Elshibli and Korpelainen, 2017; Ghazzawy *et al*., 2022).

6.5 Agroforestry and Landscape Integration

Agroforestry practices that integrate date palms with other crops improve farm sustainability and resilience to climate change. Intercropping date palms with legumes, vegetables, or forage crops enhances biodiversity, diversifies farmer income, and improves soil fertility through nitrogen fixation. Landscape-level integration of date palms with other tree species also provides windbreaks, moderates microclimates, and supports pollinator diversity. These systems contribute to ecosystem stability while reducing risks associated with monoculture cultivation (Nair, 2014; Khan, 2024).

7. Knowledge Gaps and Research Priorities

Despite growing evidence, several knowledge gaps remain. More localized studies are needed to understand the relationship between climate variables and date palm phenology. Yield-response models under future warming and rainfall scenarios are lacking. Screening of local cultivars for salinity and drought tolerance should be expanded. Socioeconomic research is needed to assess farmers' adaptive capacity, financial needs, and market risks. Addressing these gaps will support evidence-based policymaking

8. Future Research Directions

8.1 Molecular Characterization and Genome Sequencing

Molecular characterization and genome sequencing of Pakistani date palm varieties can provide insights into genetic diversity, stress tolerance, and breeding potential. Genomic resources would accelerate the development of climate-resilient varieties adapted to local environments (Elshibli and Korpelainen, 2017).

8.2 Conservation of Native Cultivars

The conservation of native Pakistani cultivars through gene banks and cryopreservation is critical for safeguarding biodiversity and ensuring a reservoir of genetic material for future breeding programs. These measures protect traditional varieties from threats posed by climate change and land-use change (Gambhir *et al*., 2024: FAO, 2022).

8.3 Development of Early Sex-Determination Techniques

Date palms are dioecious, and identifying the sex of plants at an early stage is a major challenge for farmers. The development of early sex-determination techniques using molecular markers would save time, reduce costs, and improve orchard management efficiency (Jatt *et al*., 2025: Ahmed *et al*., 2019).

8.4 Postharvest Technology Advancement

Postharvest losses remain a serious issue, especially under unpredictable rainfall. Research into solar drying, controlled atmosphere storage, and cold chain development can extend shelf life and improve the export quality of Pakistani dates. Advanced postharvest technologies are essential for strengthening international competitiveness (Solangi *et al*., 2014; TDAP, 2025b).

8.5 Socioeconomic Studies

Finally, socioeconomic research is required to understand the impacts of climate change on smallholder date farmers. Studies focusing on farmer perceptions, adaptive capacity, financial challenges, and market risks would guide policies that enhance resilience and equity in the sector (Arshad *et al*., 2019).

Conclusion

Climate change is exerting significant stress on date palm cultivation in Pakistan through rising temperatures, erratic monsoon rains, drought, salinity, and floods. These factors reduce yield, fruit quality, and farmer income, particularly in Sindh and Balochistan where production is concentrated. However, adaptation measures such as

efficient irrigation, soil conservation, integrated pest management, resilient varieties, and improved postharvest technologies offer pathways to sustain productivity. Strengthened research, farmer awareness, and supportive policies are essential to safeguard this vital crop and ensure its long-term sustainability under changing climatic conditions.

Conflict of Interest

Author declares there is no any conflict of interest.

REFERENCES

Abul-Soad, A. A. (2010). *Date palm in Pakistan, current status and prospective*. USAID Firms project, p.9-11. https://doi.org/10.17660/ActaHortic.2010.882.8

Abul-Soad, A. A., S. M. Mahdi and G. S. Markhand (2015). Date palm status and perspective in Pakistan. In: *Date Palm Genetic Resources and Utilization: Volume 2: Asia and Europe* (pp. 153-205). Dordrecht: Springer Netherlands.

Ahmed, T., M. Shahid and S. Malik (2019). Molecular markers for early sex determination in date palm (*Phoenix dactylifera* L.). *Journal of Plant Breeding and Genetics*, 7(2): 45–52.

Arshad, M., H. Kächele, T. J. Krupnik, T. S. Amjath-Babu, S. Aravindakshan, A. Abbas and K. Müller (2017). Climate variability, farmland value, and farmers' perceptions of climate change: implications for adaptation in rural Pakistan. *International Journal of Sustainable Development and World Ecology*, 24(6): 532-544.

Bentrad, N. and A. Hamida-Ferhat (2020). Date palm fruit (Phoenix dactylifera): Nutritional values and potential benefits on health. In: *The Mediterranean Diet* (pp. 239-255). Academic Press.

Dawn. (2022, August 3). Rain-driven date destruction. Dawn. https://www.dawn.com/news/1702644

Elshibli, S. and H. Korpelainen (2017). Advances in molecular breeding of date palm. *Tree Genetics and Genomes*, 13(3): 1–13.

FAO. (2022). *A rapid geospatial flood impact assessment in Pakistan* (August 2022). Food and Agriculture Organization. https://www.fao.org

FAO. (2022). FAOSTAT: *Date palm production statistics*. Food and Agriculture Organization. https://www.fao.org/faostat

FAO. (2022). *Salinity and water management challenges in Pakistan's agriculture*. Food and Agriculture Organization Report.

Gambhir, M. R., S. Lal, S. H. Shaikh, S. A. S. Zaidi, F. A. Khuhro, A. R. Kanhar, ……. and M. Ali (2025). Unsafe sips: prevalence of pathogenic bacterial diversity in drinking water of Khairpur city Pakistan. *Insights-Journal of Health and Rehabilitation*, *3*(1 (Health and Allied)): 141-148.

Ghazzawy, H. S., A. E. E. Sobaih and H. A. Mansour (2022). The role of micro-irrigation systems in date palm production and quality. *Agriculture*, 12(12): 2018. https://doi.org/10.3390/agriculture12122018

Hussain, A., S. Rehman and N. Akhtar (2021). Soil conservation techniques in arid agro-ecosystems of Pakistan. *International Soil Conservation Journal*, 29(1): 55–63.

IPCC. (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability*. Cambridge University Press. https://www.ipcc.ch/report/ar6/wg2/

Jatt, T., M. S. Lee, A. L. Rayburn, M. A. Jatoi and A. A. Mirani (2019). Determination of genome size variations among different date palm cultivars (Phoenix dactylifera L.) by flow cytometry. *3 Biotech*, 9(12): 457

Jatt, T., M. A. Maitlo, S. H. Shaikh and D. P. Z. Catanho (2023). Application of Flow Cytometry in Biological Sciences. In: *Genome Size and Genetic Homogeneity of Regenerated Plants: Methods and Applications* (pp. 232-271). Bentham Science Publishers.

Jatt, T., G. S. Markhand, M. A. Jatoi, R. Ming, C. Deng, A. A. Mirani and N. Solangi (2025). Karyotype Evolution and Genetic Variation in Date Palm (Phoenix dactylifera L.) Varieties Cultivated in Pakistan. *Planta Animalia*, 4(2): 145-154.

Khan, G. A. (2024, August 11). *Experts call for boosting date palm industry*. The Express Tribune. https://tribune.com.pk

Maitlo, M, A., M. A. Qazi and A. H. Shar (2024a). Exploring the potential of biosurfactant-producing endophytic Pseudomonas aeruginosa SF1R in medical therapeutics. *Journal of Population Therapeutics and Clinical Pharmacology*, 31(9): 4378-4386. https://doi.org/10.53555/herncx95

Maitlo, M. A., A. H. Phulpoto, M. A. Qazi, S. H. Shaikh, S. Lal and N. A. Kanhar (2024b). Biocatalytic Strategies for Environmental Pollutant Degradation: Advancing Enzymatic and Molecular Innovations. *Journal of Microbiological Sciences*, 3(01): 66-81.

Maitlo, M. A., M. A. M. A. Qazi, S. H. Shaikh, I. A. Phulpoto, R. R. Maitlo, M. S. I. Bhatti, ….... and A. H. Shar (2025). Microbial Biosurfactants: A review of their Classification and Multidimensional Applications. *Planta Animalia*, 4(2): 155-163.

Nair, P. R., B. M. Kumar and V. D. Nair (2022). Classification of agroforestry systems. In: *An introduction to agroforestry: four decades of scientific developments* (pp. 29-44). Cham: Springer International Publishing.

Naveed, H., V. Andoh, W. Islam, L. Chen and K. Chen (2023). Sustainable pest management in date palm ecosystems: Unveiling the ecological dynamics of Red Palm Weevil (Coleoptera: Curculionidae) infestations. *Insects*, 14(11): 859. https://doi.org/10.3390/insects14110859

Pakistan Meteorological Department (PMD). (2024). *State of Pakistan Climate in 2023*. Islamabad: PMD.

Solangi, N. (2014). Accelerated ripening of Aseel dates using sodium chloride to avoid monsoon deterioration. *In: Proceedings of the 5th International Date Palm Conference*.

Solangi, N., A. A. Abul-Soad, M. A. Jatoi, A. A. Mirani and G. S. Markhand (2024). Date Palm Cultivation, Consumption and Export: Current Status and Future Challenges-A Review. *Proceedings of the Pakistan Academy of Sciences: B. Life and Environmental Sciences*, 61(4): 333-350.

Soomro, M. H., J. M. Mari, I. A. Nizamani and A. A. Gilal (2022). Performance of Ferrolure+ pheromone in the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) management in date palm growing areas of Sindh, Pakistan. *Journal of the Saudi Society of Agricultural Sciences*, 21(2): 114-124.

TDAP. (2025a). *Dates production, consumption, exports and potential*. Trade Development Authority of Pakistan Report.

TDAP. (2025b). *Pakistan Dates Sector: Production, Processing and Export Potential*. Trade Development Authority of Pakistan.

You, Y., M. Ting and M. Biasutti (2024). Climate warming contributes to the record-shattering 2022 Pakistan rainfall. *Npj, Climate and Atmospheric Science*, 7(1): 89.

(*Accepted for publication October 2025*)