

NATURAL DIET AND FEEDING HABITS OF THE *SARDINELLA LONGICEPS* VAL. (OIL SARDINE) OF BALOCHISTAN COAST, PAKISTAN

Farzana Yousuf, Naeema Elahi and Sadaf Tabassum

Department of Zoology, University of Karachi, Karachi-75270, Pakistan.

ABSTRACT

The food and feeding habits of *Sardinella longiceps* Val. (oil sardine), an important food item from coast of Gwadar, Balochistan was investigated for 12 months period. Four hundred and thirty specimens of *S. longiceps* Val. were examined for food and feeding habits. It was caught by gear operating in the sea floor. Fish obtained from the surface gill net showed poor feeding and the food consisted mainly of *Coscinodiscus*, a typical diatom. The fish from bottom gill nets and trawl nets had gorged stomachs and the food item comprised mainly, either *Pleurosigma* or *Ornithocercus*, typical bottom-dwelling forms. There was variation in feeding in relation to size.

Key- words: *Sardinella longiceps* Val., Food, Feeding habits, Balochistan coast, Marine water fishes.

INTRODUCTION

The oil sardine, *Sardinella longiceps* Val. is a very important pelagic fish. Besides, being a favored nutritionally rich and affordable table fish occurring abundantly almost throughout the year, it also serves as a source for valuable by-products like sardine oil used in several industries and fish-meal for cattle and poultry feed production. Its fishery is characterized by remarkably wide fluctuations on a seasonal, annual and decadal scale. The successful years of oil sardine fishery bring as much prosperity to the fishing community as its failure a major economic setback.

The biology of Oil sardine grows rapidly during the first few months and matures early within its life span of about 2 ½ years. The age at first maturity occurs at less than one year, at about 150 mm size. Maturation is controlled by climatic factors like temperature and intensity of rainfall. The oil sardine is a planktivore and diatoms, dinoflagellates and copepods are the favoured food items. The abundance of diatom is said to indicate abundance of oil sardine in coastal waters. The optimum temperature and salinity ranges for distribution and abundance of oil sardine is 27-28°C and 22.8-33.5 ppt, respectively. The food and feeding habits of *Sardinella longiceps* Val., economically the most valuable clupeiform fish from, had been the subject of study for the past five decades (Hornell, 1910; Homell and Nayudu, 1924; Devanesan, 1943; Chidarbaram, 1950; Nair, 1953; Nair and Subrahmanyam, 1955; Venkataraman, 1960; Dhulkhed, 1962). Although a general idea of the elements on which the oil sardine subsists is worked out, data published on the food-habits of the different stages in its life-span is scanty and doubtful. Chidambaram (1950) opines that the food of young and adult sardines does not differ much.

Reference that "immature adults and spawners" feed "mostly on phytoplankton" and "very young sardines" feed mainly on copepods has been made by Nair (1953). Similar feeding habits have been reported for other fishes by Abdel Aziz and Gharib (2007), Joadder (2007), Kahilainen (2004), Kumar and Balasubrahmanyam (1989), Nunn *et al.*, (2007) (Oronsaye (2005), Shalloof (2009), Stergiou and Karpouzi (2002) and Tsiklirasi (2005).

The aims of this study were to analyze the natural diet and feeding habits of *S. longiceps* Val. inhabiting Gwadar, Balochistan coast.

MATERIALS AND METHODS

Gwadar is situated towards the western end of Pakistan's Makran coast, some 40 km east of the border with Iran. Gwadar is a rocky headland connected to the mainland by a sandspit. The headland stretches east-west for 13 km, with a maximum width of 3 km. The 430 specimens of *Sardinella longiceps* Val. were obtained from Gwadar coast from January 2009 to December 2009. The fish were transferred to a deep freezer. The preserved specimens were dry before analysis. The standard and total length (in mm) were measured on a measuring board while the weights (in grams) were determined using a balance. The stomach contents were emptied into a Petri dish and examined under a binocular microscope. The state of fullness of stomach was recorded. The volume of stomach content was determined by the displacement method. In cases where the stomach gorged, the gut content was made up to a known volume (100 ml) and whenever the food was found to be dense, a sub-sample of this was taken and further diluted. An aliquot of 1 ml was taken and spread over a counting chamber. The number method (Hynes, 1950) was followed for the estimation of relative items of food was calculated in terms of percentages.

RESULTS AND DISCUSSION

Four hundred and thirty specimens of *Sardinella longiceps* Val. were examined for food and feeding habits. Two hundred and fourteen fishes (49.76 %) had empty stomach (Table 1). Analysis of monthly variation in empty stomach (Table 1) showed that the highest occurrence of empty stomachs was in August 2009 (80%), while the lowest was noted in February and May 2009 (25%). Variation in empty stomach by size group (Table 2) indicated that the small size group of *Sardinella longiceps* Val. had the least number of empty stomachs (50.84 %). The stomach contents of *S. longiceps* Val. are presented in Table 3 and the summary of the items is illustrated in Fig 1. The fish had fed almost exclusively on a dinoflagellate, *Ornithocercus*. Zooplankters (copepods) were noticed to a negligible extent in the diet almost all around the years. The difference in the gut contents of fish taken from the surface waters and close to the sea floor was evident. In fish from bottom nets the main item of food was *Pleurosigma* or *Ornithocercus*, which are typical bottom dwelling forms.

In the gut of fish from surface gill net on the other hand, *Coscinodiscus*, a typical diatom of the upper layers of water, formed an important item of food. Thus, it appeared that fish collected from different levels have been feeding on the different species of diatoms or dinoflagellates which were dominant in the respective layers of water. The occurrence of *Pleurosigma* or *Ornithocercus* in the bottom layers may be due to the decrease in the viscosity of surface waters consequent on the increase in temperature and also to the condition of optimum light (Subrahmanyam, 1959; Raymount, 1963). The intensity of grazing as evidenced by the occurrence of single species of phytoplankton in the stomach of oil sardine caught in 10-20 m depth indicates that very rich feeding grounds exist at these depths. The observations also indicate that the oil sardine occasionally resorts to a demersal habitat.

Table 1. Monthly variation in empty stomach of *Sardinella longiceps* Val. from Balochistan coast.

Month (2009)	Number of fish examined	Number with empty stomachs	% Empty stomachs
January	30	10	33.33
February	20	05	25
March	35	20	57.14
April	35	15	43.85
May	20	05	2.5
June	30	20	66.66
July	30	10	33.33
August	50	40	80
September	50	30	60
October	50	20	40
November	30	14	46.66
December	50	25	50
TOTAL	430	214	49.76%

Table 2. Variation in empty stomach by size of *Sardinella longiceps* Val. from Balochistan coast.

Size/Standard length mm	Number of fish examined	Number with empty stomachs	% Empty stomachs
Small-sized fish (155-175)	141	60	42.53
Medium-sized fish (180-200)	171	85	49.7
Large-sized fish (205-220)	118	60	50.84

The other workers, Hornell and Nayudu (1924) stated that the oil sardine "feed upon immense quantities of unicellular plants (Phytoplankton) and animals". John and Menon (1942), Chidambaram (1950) and Nair (1953) observed that diatoms, dinoflagellates and copepods constituted food of oil sardine. It is to be noted here that these workers carried out their investigations at Gwadar. The present study has shown that in the Gwadar waters also the commercial size groups of oil sardine feed entirely on plankton, especially on phytoplankton.

Nair and Subrahmanyam (*op. cit*) have stated that the diatom is the favourite food item of oil sardine at Gwadar. Other important species of diatoms were *Coscinodiscus*, *Biddulphia* and *Pleurosigma*. Dinoflagellates were also seen. A large numbers on a number of copepods were also observed in the gut inclusion but in lesser numbers.

Table 3. Various food items and displacement volume of stomach of *Sardinella longiceps* Val.

No. of specimen examined	Size range of oil sardine (mm)	Displacement value of stomach (ml) range	Pleuro-sigma	Conscinoduscus	Biddulpnia	Diplonies	Ornithocercus	Dinophysis	Copepods
141	155-175	1.1-2.0	38.46	21.79	7.69	12.82	5.12	7.69	6.4
171	180-200	0.8-2.5	36.36	27.2	0.90	3.63	9.09	4.54	18.18
118	205-220	0.4-2.9	24.39	12.19	2.43	7.319	12.19	4.87	36.58




Fig. 1. Stomach contents of *Sardinella longiceps* Val., from Balochistan coast.

REFERENCES

Abdel Aziz, N. and S. Gharib (2007). Food and feeding habits of round sardinella (*Sardinella aurita*) in El Mex Bay, Egypt. *Egyptian Journal of Aquatic Research*, 33 (2): 202-221.

Chidambaram, K. (1950). Studies on the length-frequency of the oil sardine *Sardinella longiceps* Cuv. and Val. and on certain factors influencing their appearance on the Calicut coast of Madras Presidency. *Proc. Indian Acad. Sci.*, 31 (5): 252-286.

Devanesan, D.W. (1943). A brief investigation into the causes of the fluctuation of the annual fishery of the oil sardine of Malabar, *Sardinella longiceps* Cuv. and Val.; determination of its age and spawning ground. *Madras Fish. Bull.*, 28(1): 1-38.

Dhulkhed, M.H. (1962). Observations on the food and feeding habits of the Indian oil Sardine, *Sardinella longiceps* Val. *Indian J. Fish.*, 9 (1): 37-47

Hornell, J. (1910). Report on the results of a fishery cruise along the Malabar coast and to the Laccadive Islands in 190: *Madras Fish Bull.*, 4 (4): 71-126.

Hornell and M.R. Nayudu (1924). A contribution to the life-history of the Indian oil sardine with notes on the plankton of the Malabar coast. *Ibid.* 17 (5): 129-197.

Hynes, H. B. N. (1950). The food of the fresh water sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. *Anim. Ecol.*, 19(1): 35-58.

Joadder, Md. A. R. (2007). Food and feeding habits of *Gagata youssoufi* (Rahman) from the river Padma in Rajshahi. *Univ. J. Zool., Rajshahi University*, 25: 69-71.

John, G. G. and M. A. S. Menon (1942). Food and feeding habits of oil sardine and mackerel. *Curr.Sci.*, 11(6): 243.

Kahilainen, K. (2004). *Ecology of sympatric whitefish (Coregonus lavaretus (L.)) forms in a subarctic lake*. Ph.D. Biology dissertation, Univ. of Helsinki, Helsinki, Finland.

- Kumar, K. and K. Balasubrahmanyam (1989). Fishery and biology of oil sardine, *Sardinella longiceps*, from coastal water of India. *Central Marine Fisheries Research Institute*, 44: 42-45.
- Nair, R.V. (1953). Studies on the revival of the Indian Oil sardine fishery. *Proc. Indo-Pac. Fish. Counc.* 1953: 115-129.
- Nair and R. Subrahmanyam (1955). The diatom *Fragilaria oceanica* Cleve, an indicator of abundance of the Indian oil sardine, *Sardinella longiceps*. Cuv. and Val. *Curr. Sci.*, 24: 41-42.
- Nunn, A.D., J.P. Harvey and I. G. Cowx (2007). The food and feeding relationships of larval and 0+ year juvenile fishes in lowland rivers and connected waterbodies. II. Preyselection and the influence of gape. *J. Fish Biol.*, 70: 743-757.
- Oronsaye, C.G. and F.A. Nakpodia (2005). A comparative study of the food and feeding habits of *Chrysichthys nigrodigitatus* and *Brycinus nurse* in a tropical river. *Pak. J. Sci. Ind. Res.*, 48: 118-121.
- Raymont, J. F. G. (1963). *Plankton arui productivity in the oceans*. Pergamon Press.
- Shalloof, K.A. and N. Khalifa (2009). Stomach contents and feeding habits of *Oreochromis niloticus* (L.) from Abu-Zabal lakes, Egypt. *World Applied Sciences Journal*, 6 (1): 1-5.
- Stergiou, K.I. and V.S. Karpouzi (2002). Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries*, 11: 217-254.
- Subrahmanyam, R. (1959). Studies on the phytoplankton of the west coast of India. Part I. *Proc. Indian Acad. Sci.*, 50B(3): 113-187.
- Tsiklirasi, A.C., M. Torre and K. Stergiou (2005). Feeding habits and trophic level of round sardinella (*Sardinella aurita*) in the northeastern Mediterranean (Aegean Sea, Greece). *Journal of Biological Research*, 3: 67-75.
- Venkataraman, G. (1960). Studies on the food and feeding relationships of the inshore fishes off Calicut in the Malabar coast. *Ibid.*, 7 (2): 275-306.

(Accepted for publication May 2012)