

DISTRIBUTION OF EPIPHYTIC CYANOBACTERIA ON GREEN MACROALGAL SPECIES OCCURRING AT A ROCKY SHORE (BULEJI), KARACHI, PAKISTAN

Azra Bano¹ and Pirzada. J. A. Siddiqui²

¹Lasbela University of Agriculture Water and Marine Sciences, Uthal-90150, Baluchistan,

²Centre Of Excellence in Marine Biology, University of Karachi, Karachi-75270, Pakistan

*Corresponding author: Azra Bano; azrabano2006@yahoo.com

ABSTRACT

The present study is the part of research concerning identification of cyanobacteria from coastal water of Karachi. As report on the distribution cyanobacterial epiphytes in coastal area of Pakistan is scarce. In this study sixty eight species of cyanobacteria were identified from eight marine green macroalgal species collected from Buleji, a rocky coast of Karachi. The majority of cyanobacteria were from the order Chroococcales and Nostocales and only a small number of species were found to belong to Chamaesiphonales and Pleurocapsales. The epiphytic species consists of 27 unicellular and 41 filamentous non-heterocystous cyanobacterial species. Heterocystous forms were not observed during these studies. Cyanobacterial species exhibited a variable capability to grow in a variety of media. Also, some cyanobacteria were observed specifically associated on particular macroalgal species, while other were widespread on all species.

Key-words: Buleji, Cyanobacteria, Epiphytic, Marine habitat.

INTRODUCTION

Cyanobacteria occur in a variety of habitats of tropical, sub-tropical and temperate areas in both marine and freshwater environments (Sukenik *et al.*, 2012; Murrell and Loes, 2004; Saker and Griffiths 2001; Hoffmann, 1999; Desikachary 1959). Endophytic cyanobacteria are known from terrestrial and freshwater environments. For example, endosymbiosis relationship has been established for water fern *Azolla* (Baker, *et al.*, 2003; Stewart *et al.*, 1980; Singh and Singh, 1988). Cycads (Ow *et al.*, 1999; Peter *et al.*, 1986; Millbank, 1974), lichens (Wirtz *et al.*, 2003; Fogg *et al.*, 1973; Rai, 1990; Desikachary, 1959) and an angiosperm *Gunnera* (Bergman and Osborne, 2002; Silvester, 1976). An endophytic cyanobacteria *Richellia intracellularis* is also reported in a microalgal species *Rhizosolenia styloformis* inhabiting in Tropical Ocean. *Richellia-Rhizosolenia* association is probably the only known endosymbiosis established from marine environment. (Madhu *et al.*, 2013; Bergman *et al.*, 1992). The epiphytic cyanobacterial species on marine plants are comparatively better described. Cyanobacteria has been shown associated with seagrass, *Zostera marina* (Hamisi *et al.*, 2013; Pinckney and Micheli, 1998; Penhale, 1977; Penhale and Smith, 1977; Sand- Jansen, 1977), mangrove barks and pneumatophores (Naidoo *et al.*, 2008; Potts, 1980), macroalgae (Carpenter, 1972; Cattaeno and Kalff, 1979, 1980; Desikachary, 1959) etc. Most of these species were from terrestrial and freshwater habitats. Comparatively a little work has been done on the occurrence of cyanobacteria associated particularly with marine plants. Most epiphytic cyanobacteria are briefly mentioned in the literature. There has been no dedicated work as regard to epiphytic cyanobacteria associated with marine algae, particularly from Pakistan. Therefore, present study was undertaken to report on the species diversity of cyanobacteria occurring as epiphytes on six species of macroalgae.

MATERIALS AND METHODS

Cyanobacterial association with marine flora

A total of 8 species of seaweeds were randomly collected from Buleji, a rocky coast of Karachi. List of the species is set out in Table (1). Seaweeds species were kept separately in polythene bags and brought to the laboratory. The seaweeds were gently washed with sterilized seawater to remove loosely attached epiphytic species and other organisms. The seaweeds were divided into different portions, for example, frond, utricle and rhizoids, each portions was separately inoculated in tubes containing three different types of media ASNIII, MN and Miquel's media and incubated under cool florescent light (12L/ 12D) at 30 ± 2°C.

All seaweeds were inoculated in triplicates and in some cases where growth was either low or unobservable experiments were repeated more rigorously. The field materials and growth obtained in culture tubes were observed under the light microscope. The taxonomic assessments were done with the help of literature (Rippka *et al.*, 1979; Desikachary, 1959; Anagnostidis and Komarek, 1985, 1988, Komarek and Anagnostidis, 1986, 1989).

RESULTS

Cyanobacterial associates of macroalgal species

The study on the association of cyanobacteria with marine algae revealed that a number of species are living attached to macrophytes in nature. The study was performed on 8 green algal species (Chlorophytes) (Table 1).

However, 41 species were exclusively associated with green seaweed species. The epiphytic cyanobacteria were either unicellular form or non-heterocystous filamentous species. The heterocystous form was not observed. This research constitutes first detail report on the epiphytic cyanobacteria from Pakistani waters.

In total, 68 cyanobacteria species were identified on the green seaweeds. They were predominantly Chroococcalean (18 species) and Nostoclean (41 species) (Tables 1 and 2). The highest number of species were recorded in the genus *Phormidium* (23 species), followed in the descending order of the number of representing species by genera: *Oscillatoria* (9 species), *Chroococcus* (8 species), *Dermocarpa* (5 species), *Spirulina* (3 species), *Lyngbea* (3 species), *Merismopedia* (3 species), *Synechocystis* (2 species), *Aphanocapsa* (2 species), and *Myxosarcina* (2 species). The genera with only one representative species were *Planktothrix*, *Komvophoron*, *Pseudoanabaena*, *Gloeocapsa*, *Synchococcus*, *Gloeotheca*, *Chroococciopsis* and *Chamaecalyx* (Table 2).

Considering the individual green algal species, it may be deduce that there are some (15 species) more common cyanobacteria which associated with three or more algal species. It is also observed that group of cyanobacteria belonging to Chroococcales, Chamaesiphonales and Pleurocapsales were not associated with *Ch. Prostrata*, *E. intestinales*, *V. pachynema*, *C. taxifolia*, *C. peltata* and *U. indica* (Table 2 and 3).

Some species of cyanobacteria appeared to have associated with only one algal species, i.e., algal species-specific association. Out of 68 species, 41 species had macroalgal species-specific association most of which observed on *E. intestinales* and *C. peltata* (9 species each). *U. indica*, *U. rigida*, *Ch. Prostrate*, *Bryopsis pennata*, *V. pachynema* and *C. taxifolia* had 8, 5, 5, 4, 3, 2 and 2, species of cyanobacteria attached with them, respectively (Fig.1. and Table 1).

Three types of media (ASNIII, MN and Miquel's) were used to improve the chances of obtaining obtained in above three media media (Table 2). The highest numbers of species (40 species) were observed in Miquel's medium. Thirty six species were able to grow in ASNII medium and only 25 species were obtained from culture in MN medium (Table 1). Although some cyanobacteria (8 species) were able to grow in all three media (Table 1, Fig. 2.), there were many species which were observed in one type of medium. ASNIII, MN and Miquel's media showed 15, 11 and 17 species that grew in the media, respectively (Table 2 and Fig. 2.).




Fig. 1. Species-specific cyanobacteria associated with A- *Enteromorpha intestinalis*, B- *Ulva indica*, C- *Ulva rigida*, D- *Chaetomorpha prostrata*, E- *Valoniopsis pachynema*, F- *Bryopsis pennata*, G- *Caulerpa peltata*, H- *Caulerpa taxifolia*.




Fig. 2. Cyanobacterial species were common in all three media but several species showed up only in one specific medium.

Table 1. List of chlorophyte species studied for the epiphytic species of cyanobacteria.

Bryopsidophyceae	Ulvophyceae
<i>Caulerpa peltata</i>	<i>Enteromorpha intestinalis</i>
<i>Caulerpa taxifolia</i>	<i>Ulva indica</i>
	<i>Ulva rigida</i>
	<i>Chaetomorpha prostata</i>
	<i>Valoniopsis pachynema</i>
	<i>Bryopsis pennata</i>

DISCUSSION

The data presented here contributes to the regional diversity data pool, and at the same time confirms the presence of many species cyanobacteria reported from the other part of the world in the northern Arabian sea bordering Pakistan.

Only a few reports are available on epiphytic cyanobacteria from coastal zones of Pakistan. These study reports only a few species of macroalgae (Nizamuddin and Gessner, 1970; Shameel, 1987; Bano and Siddiqui, 2003) and on mangrove plants (Saifullah and Taj, 1995; Siddiqui and Bano, 2001).

Although its importance in the aquatic ecosystem, marine cyanobacteria has not been studied in detail. The growth of cyanobacteria on the group of macroalgae may suggest its possible role in the physiology of the algal species (Sellner, 1997; Philips and Zeman, 1990). The epiphytic cyanobacterial relationship may be symbiotic relationship. It has previously been reported that an exchange of carbon and nitrogen compounds exist between epiphytic cyanobacteria and the marine plants (Mohamed and Shehri, 2010; Vahtera, *et al.*, 2007; [Murakami et al.](#), 2004; Toledo *et al.*, 1995; Moisander *et al.*, 2005; Harlin and Craigie, 1975; Wetzel and Penhale, 1979; Penhale and Thayer, 1980; Harlin, 1973; McRoy and Goring 1974). For example, diazotrophic epiphytes have been reported from a few marine algae (Gómez, *et al.*, 2005; Toledo *et al.*, 1995; Capone, 1977; Capone *et al.*, 1977; Carpenter, 1972; Hansen, 1977; Head and Carpenter, 1975).

Some cyanobacterial species encountered in the study as epiphytes are also present in many other niches of the rocky shore and are also reported from other habitats as well (Desikachary, 1959; Thajuddin and Subramanian, 1992, 1994; Santra and Pal 1988; Santra *et al.*, 1988). These organisms therefore, appeared to be cosmopolitan and

marine plant surface may simply be a substrate for attachment (Sukenik *et al.*, 2012; Verspagen *et al.*, 2005; Furey, 2003). However, there are some species of cyanobacteria found exclusively on macroalgae, and hence suggested a possibility of the existence of a symbiotic relationship (Lesser *et al.*, 2004). Some variation in the diversity of species associated with different seaweed species, suggest that there may be a specific relationship between some macroalgae and cyanobacteria (Thacker and Starnes, 2003; Thacker *et al.*, 2001).

Table 2. Distribution of cyanobacterial species on eight different species of chlorophytes inhabiting a rocky shore of Buleji near Karachi.

Cyanobacterial species	Media*	Macroalgal Species							
		<i>Enteromorpha intestindis</i>	<i>Ulva indica</i>	<i>Ulva rigida</i>	<i>Chaetomorpha prostata</i>	<i>Valoniopsis pachynema</i>	<i>Bryopsis pennata</i>	<i>Caulerpa peltata</i>	<i>Caulerpa taxifolia</i>
Chroococcales									
<i>Synechocystis pevalekii</i>	ASNIII, MN	-	+	+	-	-	-	+	-
<i>S. aquatilis</i>	MM	-	-	-	-	-	-	+	-
<i>Gloeocapsa cripidinum</i>	ASNIII	-	-	-	-	-	+	-	-
<i>Chroococcus cohaerence</i>	ASNIII, MN, MM	+	-	+	-	+	-	-	-
<i>C. turgidus</i>	ASNIII, MM	-	-	-	-	-	-	+	+
<i>C. minor</i>	MM	-	+	+	-	-	-	-	-
<i>C. montanus</i>	ASNIII	+	-	-	-	-	-	-	-
<i>C. tenax</i>	ASNIII, MM	-	-	-	-	-	-	+	+
<i>C. hansgirgi</i>	ASNIII	-	-	-	-	-	-	+	-
<i>C. dispersus</i>	MM	-	-	-	-	-	-	-	+
<i>C. pallidus</i>	ASNIII	+	-	-	-	-	-	-	-
<i>Synechococcus elongatus</i>	ASNIII, MM	-	+	+	-	-	-	-	+
<i>Aphanocapsa littoralis</i>	MN	-	+	-	-	-	-	-	-
<i>A. rivulais</i>	MN	-	+	-	-	-	-	-	-
<i>Gloeothece rhodochlmys</i>	ASNIII, MN	-	-	+	-	-	-	-	-
<i>Merismopedia elegans</i>	ASNIII, MN	-	+	+	-	-	+	+	-
<i>M. punctata</i>	MN	-	+	-	-	-	-	-	-
<i>M. tenuissima</i>	ASNIII	+	-	-	-	-	-	-	-
Chamaesiphonales									
<i>Chroococidiopsis indica</i>	MM	-	-	-	-	-	-	-	+
<i>Dermocarpa leibleiniaie</i>	ASNIII, MN, MM	-	+	+	+	-	+	-	-
<i>D. olivacea</i>	MM	-	-	+	-	-	-	-	-
<i>D. versicolor</i>	MN	-	+	-	-	-	-	-	-
<i>D. clavata</i>	MN	-	-	+	-	-	+	-	-
<i>D. sphaerica</i>	MN	-	-	+	-	-	-	-	-
<i>Chaemaecalyx swirrenkoi</i>	MN	-	-	+	-	-	-	-	-
Pleurocapsales									
<i>Myxosarcina spectabilis</i>	MN, MM	+	-	+	-	-	+	-	-

<i>M. burmensis</i>	ASNIII, MM	-	-	-	-	-	+	-	-	
Nostocales										
<i>Komvophoron minutum</i>	ASNIII	-	+	-	-	-	-	-	-	
<i>Psuedoanabaena catenata</i>	ASNIII, MN, MM	-	+	+	-	-	+	-	-	
<i>Lyngbea nordgardhii</i>	ASNIII, MM	+	-	-	-	-	-	-	-	
<i>L. lagerheimi</i>	MN	-	+	-	-	-	-	-	-	
<i>L. chlorospira</i>	ASNIII, MM	-	+	-	-	+	-	-	-	
<i>Oscillatoria limnetica</i>	ASNIII, MM	+	-	-	-	-	-	-	-	
<i>O. schultzei</i>	MM	-	-	-	+	-	-	-	-	
<i>O. freyia</i>	MM	+	-	-	-	-	-	-	-	
<i>O. guttulata</i>	MM	+	-	-	-	-	-	-	-	
<i>O. splendida</i>	MM	+	-	-	-	-	-	-	-	
<i>O. pseudogaminata</i>	ASNIII, MN, MM	-	+	+	+	+	+	-	-	
<i>O. wellei</i>	MN	-	-	-	+	-	-	-	-	
<i>O. proteus</i>	ASNIII	-	-	-	+	-	-	-	-	
<i>O. deflexa</i>	MM	-	-	-	+	-	-	-	-	
<i>Spirulina labyrinthiformis</i>	ASNIII, MN, MM	+	+	-	-	-	+	+	-	
<i>S. subsalsa</i>	ASNIII	-	-	-	-	-	-	+	-	
<i>S. major</i>	MN	-	-	-	-	-	-	+	-	
<i>Planktothrix clathrata</i>	MM	+	-	-	-	-	-	-	-	
<i>Phormidium purpurascence</i>	ASNIII, MN, MM	-	-	-	-	-	+	+	+	
<i>P. laminosum</i>	ASNIII, MM	-	-	-	-	+	-	+	+	
<i>P. tenue</i>	ASNIII, MN, MM	-	-	-	-	+	+	+	-	
<i>P. insigne</i>	ASNIII	-	-	+	+	-	-	-	-	
<i>P. angustissimum</i>	ASNIII, MM	-	-	-	+	+	-	+	-	
<i>P. acutissimum</i>	MM	-	-	-	-	+	-	-	-	
<i>P. ambiguum</i>	ASNIII	-	-	-	+	-	-	+	-	
<i>P. africanum</i>	MN, MM	-	+	-	-	-	-	+	-	
<i>P. amplivaginatum</i>	MM	-	-	-	-	+	-	-	-	
<i>P. fragile</i>	ASNIII, MN, MM	+	-	+	-	+	+	-	-	
<i>P. inudatum</i>	ASNIII, MM	-	-	-	-	-	-	+	-	
<i>P. incrustatum</i>	ASNIII	-	+	-	-	-	-	-	-	
<i>P. retzii</i>	MM	-	-	-	-	-	-	+	+	
<i>P. mucicola</i>	MM	-	-	+	-	-	-	-	-	
<i>P. luteum</i>	ASNIII	-	+	-	-	-	-	-	-	
<i>P. papyraceum</i>	MN, MM	-	-	-	-	+	+	-	-	
<i>P. valdriianum</i>	MN, MM	-	-	+	-	+	-	+	-	
<i>P. jadianianum</i>	MM	-	-	-	-	-	-	+	-	
<i>P. subincrustedum</i>	ASNIII	-	-	-	-	-	-	+	-	
<i>P. anamola</i>	ASNIII	-	-	-	-	-	-	+	-	
<i>P. luridum</i>	ASNIII	-	-	-	-	-	-	+	-	
<i>P. laetevirens</i>	MM	-	-	-	+	-	-	-	-	
<i>P. rimosum</i>	MN	-	-	-	-	-	+	-	-	

*ASNIII, MN (Rippka et al 1979), MM (Imai 1977)

Table 3. Total number of species observed on surface of chlorophyte species. Species are also grouped according to their taxonomic orders.

Cyanobacterial orders	<i>Enteromorpha intestinalis</i>	<i>Ulva indica</i>	<i>Ulva rigida</i>	<i>Chaetomorpha prostate</i>	<i>Valoniopsis pachynema</i>	<i>Bryopsis pennata</i>	<i>Caulerpa peltata</i>	<i>Caulerpa taxifolia</i>
Chroococcales	4	7	6	-	1	2	6	4
Chamaesiphonales	-	2	5	1	-	2	-	1
Pleurocapsales	1	-	1	-	-	2	-	-
Nostocales	8	9	6	9	10	8	16	3
Total	13	18	18	10	11	14	22	8

ACKNOWLEDGMENTS

The authors are thankful to Vice Chancellor, Lasbela University of Agriculture Water and Marine Science, for his invariable support for providing facility. We also thank to staff of faculty of Marine Science of LUAWMS for their assistance during preparation of this manuscript.

REFERENCES

- Anagnostidis, K. and J. Komárek (1985). Modern approach to the classification system of cyanophytes, 1– Introduction. – *Arch. Hydrobiol. Suppl. 71/Algological Studies* 38/39: 291-302.
- Anagnostidis, K. and J. Komárek (1988). Modern approach to the classification system of cyanophytes, 3– Oscillatoriales. – *Arch. Hydrobiol. 80/Algological Studies* 50–53: 327-472.
- Baker, J. A., B. Entsch and D. B. McKay (2003). The cyanobiont in an *Azolla* fern is neither *Anabaena* nor *Nostoc*. *FEMS. Microb. Lett.* 229: 43-47.
- Bano, A. and P. J. A. Siddiqui (2003). Intertidal cyanobacterial diversity on a rocky shore at Buleji near Karachi, Pakistan. *Pakistan J. Bot.*, 35 (1): 27-36.
- Bergman, B. and B. Osborne (2002). The *Gunnera*: *Nostoc* Symbiosis. Biology and Environment: *Proceedings of the Royal Irish Academy*. 102B(1): 35-39.
- Bergman, B., C. Johansson and E. Söderbäck (1992). The *Nostoc*–*Gunnera* Symbiosis. *New Phytologist*. 122: 379-400.
- Capone, D. G. (1977). N₂ (C₂H₂) fixation by macroalgal epiphytes. In *Proc. 3rd int. coral Reef Symp* [Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Florida] (pp. 337-342).
- Capone, D. G., D. L. Taylor and B. F. Taylor (1977). Nitrogen fixation (acetylene reduction) associated with macroalgae in a coral-reef community in the Bahamas. *Marine Biology*. 40 (1): 29-32.
- Carpenter, E.J. (1972). Nitrogen fixation by blue-green epiphytes on pelagic *Sargassum*. *Science*. 178: 1207-1209.
- Cattaeno, A. and J. Kalf (1979). Primary production of algae growing in natural and artificial aquatic plants. A study of interaction between epiphytes and their substrates. *Limnol. Oceanogr.*, 24 (6): 1031-1037.
- Desikachary, T.V. (1959). Cyanophyta. Monographs in algae. New Delhi. Indian council of Agricultural Research.
- Fogg, G. E., Stewart, W. D. P., Fay, P. and Walsby, A. E. (1973). *The blue-green algae*. Pp. 495. Academic Press, N. Y.
- Furey, P. C. (2003). Cosmopolitan cyanobacteria. *The American biology teacher*. 65(8): 595-598.
- Gómez, F., K. Furuya and S. Takeda (2005). Distribution of the cyanobacterium *Richelia intracellularis* as an epiphyte of the diatom *Chaetoceros compressus* in the western Pacific Ocean. *J. of Plank. Res.* 27(4): 323-330.
- Hamisi, M., B. Díez, T. Lyimo, K. Ininbergs and B. Bergman (2013). Epiphytic cyanobacteria of the seagrass *Cymodocea rotundata*: diversity, diel nifH expression and nitrogenase activity. *Environ. Microb. Rep.* 5(3): 367-376.
- Hanson, R. B. (1977). Pelagic *Sargassum* community metabolism: Carbon and nitrogen. *J. of Exp. Mar. Biol. and Ecol.* 29(2): 107-118.

- Harlin, M. M. and J. S. Craigie (1975). The distribution of photosynthate in *Ascophyllum nodosum* as it relates to epiphytic *Polysiphonia lanosa* 12. *J. of Phy.* 11(1): 109-113.
- Harlin, M. M. (1973). Transfer of products between epiphytic marine algae and host plants 1, 2. *J. of Phyco.*, 9(3): 243-248.
- Head, W. D. and E. J. Carpenete, (1975). Nitrogen fixation associated with marine macroalgae *Codium fragile*. *Limno. Oceanog.*, 20: 815-823.
- Hoffmann, L. (1999). Marine cyanobacteria in tropical regions: diversity and ecology. *Eur. J. Phycol.* 34: 371-379.
- Komárek, J. and K. Anagnostidis (1986). Modern approach to the classification system of cyanophytes, 2–Chroococcales. – *Arch. Hydrobiol. 73/Algological Studies* 43:157-226.
- Komárek, J. and K. Anagnostidis (1989). Modern approach to the classification system of Cyanophytes, 4–Nostocales. – *Arch. Hydrobiol. Suppl. 82/Algological Studies* 56: 247-345.
- Lesser, M. P., C. H. Mazel, M. Y. Gorbunov and P. G. Falkowski (2004). Discovery of symbiotic nitrogen-fixing cyanobacteria in corals. *Science*. 305(5686): 997-1000.
- Madhu, N. V., Meenu Paul, N. Ullas, R. Ashwini and T. V. Rehitha (2013). Occurrence of cyanobacteria (*Richelia intracellularis*)-diatom (*Rhizosolenia hebetata*) consortium in the Palk Bay, Southeast coast of India. *Indian J. Geo-Marine Sciences*, 42(4):453-457.
- McRoy, C. P. and J. J. Goering (1974). Nutrient transfer between the seagrass *Zostera marina* and its epiphytes. *Nature*. 248: 173-174.
- Millbank, J. W. (1974). In: *The Biology of Nitrogen fixation* (A Quispel, ed.), pp. 238-264. North Holland, Amsterdam.
- Moisander, P. H., M. F. Piehler and H. W. Paerl (2005). Diversity and activity of epiphytic nitrogen-fixers on standing dead stems of the salt marsh grass *Spartina alterniflora*. *Aquatic Microbial Ecology*. 39(3): 271-279.
- Mohamed, Z. A. and A. A. Shehri (2010). Microcystin production in epiphytic cyanobacteria on submerged macrophytes. *Toxicon*. 55, (7): 1346-1352.
- Murakami, A., H. Miyashita, M. Iseki, K. Adachi and M. Mimuro (2004). Chlorophyll *d* in an epiphytic cyanobacterium of red algae. *Science*. 303 (5664): 1633.
- Murrell, M. C. and E. M. Lores (2004). Phytoplankton and zooplankton seasonal dynamics in a subtropical estuary: importance of cyanobacteria. *Journal of Plankton Research*. 26 (3): 371-382.
- Naidoo, Y., T. D. Steinke, F. D. Mann, A. Bhatt and S. Gairola (2008). Epiphytic organisms on the pneumatophores of the mangrove *Avicennia marina*: occurrence and possible function. *African Journal of Plant Science*. 1 (2): 012-015.
- Nizamuddin, M. and F. Gessner (1970). *The marine algae of the northern part of the Arabian Sea and of the Persian Gulf*. Gebrueder Borntraeger.
- Ow, M. C., M. Gantar and J. Elhai (1999). Reconstitution of a cycad-cyanobacterial association. *Symbiosis-Rehovot*. 27(2): 125-134.
- Penhale, P. A. and G. W. Thayer (1980). Uptake and transfer of carbon and phosphorus by eelgrass (*Zostera marina* L.) and its epiphytes. *J. of Exp. Mar. Biol. and Ecol.* 42(2): 113-123.
- Penhale, P. A. and W. O. Smith Jr. (1977). Excretion of dissolved organic carbon by eelgrass (*Zostera marina*) and its epiphytes. *Limnol. Oceanog.* 22 (3): 400-407.
- Penhale P. A. (1977). Macrophyte-epiphyte biomass and productivity in an eelgrass (*Zostera marina* L.) community. *J. Exp. Mar. Biol. and Ecol.* 26(2): 211-224.
- Peters, G. A., R. E. Toia Jr, H. E. Calvert and B. H. Marsh (1986). Lichens to *Gunnera*—with emphasis on *Azolla*. In: *Nitrogen Fixation with Non-Legumes* (pp. 17-34). Springer Netherlands.
- Pinckney, J. L. and F. Micheli (1998). Microalgae on seagrass mimics: Does epiphyte community structure differ from live seagrasses?. *J. of Exp. Mar. Biol. and Ecol.* 221(1): 59-70.
- Phlips, E. and C. Zeman (1990). Photosynthesis, growth and nitrogen fixation by epiphytic forms of filamentous cyanobacteria from pelagic *Sargassum*. *Bulletin of Marine Science*. 47 (3): 613-621.
- Potts, M. (1980). Blue-green algae (Cyanophyta) in marine coastal environments of the Sinai Peninsula; distribution, zonation, stratification and taxonomic diversity *Phycologia* 19. (1): 60-73.
- Rai, A. N. (1990). *Handbook of symbiotic cyanobacteria*. CRC Press, Inc.
- Rippka, R., J. Deruelles, J.B. Waterbury, M. Herdman and R.Y. Stanier (1979). Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* 111: 1-61.
- Saker, M. L. and D. J. Griffiths (2001). Occurrence of blooms of the cyanobacterium *Cylindrospermopsis raciborskii* (Woloszynska) Seenayya and Subba Raju in a north Queensland domestic water supply. *Mar. and Fresh. Res.* 52 (6): 907-915.
- Sand-Jensen, K. A. J. (1977). Effect of epiphytes on eelgrass photosynthesis. *Aquatic Botany*. 3: 55-63.

- Saifullah, S. M. and G. Taj (1995). Marine algal epiphytes on pneumatophores of mangroves of Karachi. *The Arabian Sea Living Marine Resources and the Environment*, AIBS, Vangurad Books (PVT) Ltd., Lahore, 407-417.
- Shameel, M. (1987). A preliminary survey of seaweeds from the coast of Lasbela, Pakistan. *Botanica Marina*, 30(6): 511-516.
- Siddiqui, P. J. A. and A. Bano (2001). Distribution of epiphytic cyanobacteria on macroalgae occurring at a rocky shore (Buleji), Karachi. *Pak. J. Mar. Biol.* 7 (1& 2): 251-259.
- Silvester, W. B. (1976). Endophyte adaptation in *Gunnera-Nostoc* symbiosis. Symbiotic nitrogen fixation in plants. Cambridge University Press: Cambridge etc. 521-37.
- Singh, R. P. and P. K. Singh (1988). Symbiotic algal nitrogenase activity and heterocyst frequency in seven *Azolla* species after phosphorus fertilization. *Hydrobiologia*, 169 (3): 313-318.
- Stewart, W. D. P. (1980). Some aspects of structure and function in n fixing cyanobacteria. *Annual Reviews in Microbiology*, 34(1): 497-536.
- Sukenik, A., O. Hadas, A. Kaplan and A. Quesada (2012). Invasion of Nostocales (cyanobacteria) to subtropical and temperate freshwater lakes—physiological, regional, and global driving forces. *Frontiers in Microbiology*, 3: 1-9.
- Santra, S.C. and U.C. Pal (1988). Marine algae of mangrove delta region West Bengal, India: Benthic forms. *Indian Biolo.* 20(2): 30-41.
- Santra, S.C., U.C. Pal, H. Maity and G. Bandhyopadhaya (1988). Blue green algae in saline habitat of West Bengal: A systematic account. *Biol. Mem.*, 14(1): 81-108.
- Sellner, K. G. (1997). Physiology, ecology, and toxic properties of marine cyanobacteria blooms. *Limno. and Ocean.* 42(5): 1089-1104.
- Toledo, G., Y. Bashan and A. Soeldner (1995). Cyanobacteria and black mangroves in Northwestern Mexico: colonization, and diurnal and seasonal nitrogen fixation on aerial roots. *Canadian Journal of Microbiology*, 41(11): 999-1011.
- Thacker, R. W. and S. Starnes (2003). Host specificity of the symbiotic cyanobacterium *Oscillatoria spongeliae* in marine sponges, *Dysidea* spp. *Marine Biology*, 142(4): 643-648.
- Thacker, R., D. Ginsburg and V. Paul (2001). Effects of herbivore exclusion and nutrient enrichment on coral reef macroalgae and cyanobacteria. *Coral Reefs*, 19(4): 318-329.
- Thajuddin, N. and G. Subramanian (1990). Cyanobacterial phytoplanktons of the Gulf of Mannar region. *Proc. Natl. Symposium on cyanobacterial nitrogen fixation*. NFBGA, IARA, New Delhi pp. 457-463.
- Thajuddin, N. and G. Subramanian (1992). Survey of cyanobacterial flora of the Southern East coast of India. *Bot. Mar.*, 35: 305-314.
- Thajuddin, N. and G. Subramanian (1994). Marine cyanobacterial flora of South India. *Current Researchers in Plant Sciences*, 1-16.
- Vahtera, E., D. J. Conley, B. G. Gustafsson, H. Kuosa, H. Pitkänen, O.P. Savchuk and F. Wulff (2007). Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management in the Baltic Sea. *AMBIO: A Journal of the Human Environment*, 36(2): 186-194.
- Verspagen, J. M., E. O. Snelder, P. M. Visser, K. D. Joehnk, B. W. Ibelings, L. R. Mur and J. Huisman (2005). Benthic–pelagic coupling in the population dynamics of the harmful cyanobacterium *Microcystis*. *Freshwater Biology*, 50(5): 854-867.
- Wetzel, R. G., and P. A. Penhale (1979). Transport of carbon and excretion of dissolved organic carbon by leaves and roots/rhizomes in seagrasses and their epiphytes. *Aquatic Botany*, 6: 149-158.
- Wirtz, N., H. T. Lumbsch, T. G. A. Green, R. Türk, A. Pintado, L. Sancho and B. Schroeter (2003). Lichen fungi have low cyanobiont selectivity in maritime Antarctica. *New Phytologist*, 160: 177-183.

(Accepted for publication December 2016)